

DEFORMATION OF PROPER ACTIONS ON REDUCTIVE HOMOGENEOUS SPACES

FANNY KASSEL

ABSTRACT. Let G be a real reductive Lie group and H a closed reductive subgroup of G . We investigate the deformation of “standard” compact quotients of G/H , *i.e.*, of quotients of G/H by discrete subgroups Γ of G that are uniform lattices in a closed reductive subgroup L of G acting properly and cocompactly on G/H . For L of real rank 1, we prove that after a small deformation in G , such a group Γ remains discrete in G and its action on G/H remains properly discontinuous and cocompact. More generally, we prove that the properness of the action of any convex cocompact subgroup of L on G/H is preserved under small deformations, and we extend this result to reductive homogeneous spaces G/H over any local field. As an application, we obtain compact quotients of $\mathrm{SO}(2n, 2)/\mathrm{U}(n, 1)$ by Zariski-dense discrete subgroups of $\mathrm{SO}(2n, 2)$ acting properly discontinuously.

1. INTRODUCTION

Let G be a real connected reductive linear Lie group and H a closed connected reductive subgroup of G . We are interested in the compact quotients of G/H by discrete subgroups Γ of G . We ask that the action of Γ on G/H be properly discontinuous in order for the quotient $\Gamma \backslash G/H$ to be Hausdorff. This imposes strong restrictions on Γ when H is noncompact. For instance, if $\mathrm{rank}_{\mathbb{R}}(G) = \mathrm{rank}_{\mathbb{R}}(H)$, then all discrete subgroups of G acting properly discontinuously on G/H are finite: this is the Calabi-Markus phenomenon [Ko1]. Usually the action of Γ on G/H is also required to be free, so that $\Gamma \backslash G/H$ be a manifold, but this condition is not very restrictive: if Γ acts properly discontinuously and cocompactly on G/H , then it is finitely generated, hence virtually torsion-free by Selberg’s lemma [Sel].

In this paper we investigate the deformation of compact quotients $\Gamma \backslash G/H$ in the important case when Γ is “standard”, *i.e.*, when Γ is a uniform lattice in some closed reductive subgroup L of G acting properly and cocompactly on G/H . Most of our results hold for reductive homogeneous spaces over any local field, but in this introduction we first consider the real case.

1.1. Deformation of compact quotients in the real case. Let G be a real reductive linear Lie group and H a closed reductive subgroup of G . In all known examples, if G/H admits a compact quotient, then there is a closed reductive subgroup L of G that acts properly and cocompactly on G/H . For instance, $L = \mathrm{U}(n, 1)$ acts properly and transitively on the $(2n + 1)$ -dimensional anti-de Sitter space $G/H = \mathrm{SO}(2n, 2)/\mathrm{SO}(2n, 1)$ (see Section 6). Any torsion-free uniform lattice Γ of such a group L acts properly

discontinuously, freely, and cocompactly on G/H ; we will say that the corresponding compact quotient $\Gamma \backslash G/H$ is *standard*. Note that L always admits torsion-free uniform lattices by [Bo1]. Kobayashi and Yoshino conjectured that any reductive homogeneous space G/H admitting compact quotients admits standard ones ([KY], Conj. 3.3.10); this conjecture remains open.

Of course, nonstandard compact quotients may also exist: this is the case for instance for $G/H = (G_0 \times G_0)/\Delta_{G_0}$, where G_0 is any reductive Lie group and Δ_{G_0} is the diagonal of $G_0 \times G_0$ (see [Ghy], [Gol], [Ko3], [Sal]). But in general we know only standard examples. In order to construct nonstandard ones, it is natural, given a reductive subgroup L of G acting properly and cocompactly on G/H , to slightly deform torsion-free uniform lattices Γ of L in G and to see whether they remain discrete in G and their action on G/H remains proper, free, and cocompact.

Since our goal is to obtain nonstandard quotients, we are not really interested in trivial deformations of Γ , *i.e.*, in deformations by conjugation, for which the quotient $\Gamma \backslash G/H$ will remain standard. The problem therefore boils down to the case when L is reductive of real rank 1. Indeed, by classical preliminary reductions, that is, considering separately the different irreducible quasifactors of Γ and possibly passing to subgroups of finite index, we may assume that Γ is irreducible and that L has no compact factor. If L is semisimple of real rank ≥ 2 , then Margulis's superrigidity theorem implies that Γ is locally rigid in G ([Mar], Cor. IX.5.9).

We prove the following result.

Theorem 1.1. *Let G be a real reductive linear Lie group, H and L two closed reductive subgroups of G . Assume that $\text{rank}_{\mathbb{R}}(L) = 1$ and that L acts properly and cocompactly on G/H . For any torsion-free uniform lattice Γ of L , there is a neighborhood $\mathcal{U} \subset \text{Hom}(\Gamma, G)$ of the natural inclusion such that any $\varphi \in \mathcal{U}$ is injective and $\varphi(\Gamma)$ is discrete in G , acting properly discontinuously and cocompactly on G/H .*

We denote by $\text{Hom}(\Gamma, G)$ the set of group homomorphisms from Γ to G , endowed with the compact-open topology. In the real case, the fact that $\varphi(\Gamma)$ remains discrete in G for $\varphi \in \text{Hom}(\Gamma, G)$ close to the natural inclusion is a general result of Guichard ([Gui], Th. 2).

Theorem 1.1 improves a result of Kobayashi ([Ko3], Th. 2.4), who considered particular homomorphisms of the form $\gamma \mapsto \gamma\psi(\gamma)$, where $\psi : \Gamma \rightarrow Z_G(L)$ is a homomorphism with values in the centralizer of L in G .

By [KY], Cor. 3.3.7, Theorem 1.1 applies to the following triples (G, H, L) :

- (1) $(\text{SO}(2n, 2), \text{SO}(2n, 1), \text{U}(n, 1))$ for $n \geq 1$,
- (2) $(\text{SO}(2n, 2), \text{U}(n, 1), \text{SO}(2n, 1))$ for $n \geq 1$,
- (3) $(\text{U}(2n, 2), \text{Sp}(n, 1), \text{U}(2n, 1))$ for $n \geq 1$,
- (4) $(\text{SO}(8, 8), \text{SO}(8, 7), \text{Spin}(8, 1))$,
- (5) $(\text{SO}(8, \mathbb{C}), \text{SO}(7, \mathbb{C}), \text{Spin}(7, 1))$,
- (6) $(\text{SO}^*(8), \text{U}(1, 3), \text{Spin}(6, 1))$,
- (7) $(\text{SO}^*(8), \text{Spin}(6, 1), \text{U}(1, 3))$,
- (8) $(\text{SO}^*(8), \text{SO}^*(6) \times \text{SO}^*(2), \text{Spin}(6, 1))$,
- (9) $(\text{SO}(4, 4), \text{Spin}(4, 3), \text{SO}(4, 1))$,
- (10) $(\text{SO}(4, 3), \text{G}_{2(2)}, \text{SO}(4, 1))$.

As mentioned above, our aim is to deform standard compact quotients of G/H into nonstandard ones, which are in some sense more generic. The best that we may hope for is to obtain Zariski-dense discrete subgroups of G acting properly discontinuously, freely, and cocompactly on G/H . Of course, even when L has real rank 1, nontrivial deformations in G of uniform lattices Γ of L do not always exist. For instance, if L is semisimple, noncompact, with no quasisimple factor locally isomorphic to $\mathrm{SO}(n, 1)$ or $\mathrm{SU}(n, 1)$, then the first cohomology group $H^1(\Gamma, \mathfrak{g})$ vanishes by [Rag], Th. 1. This, together with [Wei], implies that Γ is locally rigid in G . (Here $\mathfrak{g}$ denotes the Lie algebra of G .)

For $(G, H, L) = (\mathrm{SO}(2n, 2), \mathrm{SO}(2n, 1), \mathrm{U}(n, 1))$ with $n \geq 2$, uniform lattices Γ of L are not locally rigid in G , but a small deformation of Γ will never provide a Zariski-dense subgroup of G . Indeed, by [Rag] and [Wei] there is a neighborhood in $\mathrm{Hom}(\Gamma, G)$ of the natural inclusion whose elements are all homomorphisms of the form $\gamma \mapsto \gamma\psi(\gamma)$, where $\psi : \Gamma \rightarrow \mathrm{SO}(2n, 2)$ is a homomorphism with values in the center of $\mathrm{U}(n, 1)$.

On the other hand, for $(G, H, L) = (\mathrm{SO}(2n, 2), \mathrm{U}(n, 1), \mathrm{SO}(2n, 1))$ with $n \geq 1$, there do exist small Zariski-dense deformations of certain uniform lattices of L in G (see Section 6): such deformations can be obtained by a bending construction due to Johnson and Millson [JM]. Theorem 1.1 therefore implies the following result on the compact quotients of the homogeneous space $G/H = \mathrm{SO}(2n, 2)/\mathrm{U}(n, 1)$.

Corollary 1.2. *For any $n \geq 1$, there is a Zariski-dense discrete subgroup of $\mathrm{SO}(2n, 2)$ acting properly discontinuously, freely, and cocompactly on $\mathrm{SO}(2n, 2)/\mathrm{U}(n, 1)$.*

Note that by [KY], Prop. 3.2.7, the homogeneous space $\mathrm{SO}(2n, 2)/\mathrm{U}(n, 1)$ is a pseudo-Riemannian symmetric space of signature $(2n, n^2 - 1)$.

The existence of compact quotients of reductive homogeneous spaces by Zariski-dense discrete subgroups was known so far only for homogeneous spaces of the form $(G_0 \times G_0)/\Delta_{G_0}$.

1.2. Deformation of properly discontinuous actions over a general local field. We prove that the properness of the action is preserved under small deformations not only for real groups, but more generally for algebraic groups over any local field $\mathbf{k}$. By a local field we mean $\mathbb{R}$, $\mathbb{C}$, a finite extension of $\mathbb{Q}_p$, or the field $\mathbb{F}_q((t))$ of formal Laurent series over a finite field $\mathbb{F}_q$. Moreover we relax the assumption that Γ is a torsion-free uniform lattice of L , in the following way.

Theorem 1.3. *Let $\mathbf{k}$ be a local field, G the set of $\mathbf{k}$ -points of a reductive algebraic $\mathbf{k}$ -group $\mathbf{G}$, and H (resp. L) the set of $\mathbf{k}$ -points of a closed reductive subgroup $\mathbf{H}$ (resp. $\mathbf{L}$) of $\mathbf{G}$. Assume that $\mathrm{rank}_{\mathbf{k}}(\mathbf{L}) = 1$ and that L acts properly on G/H . If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, let Γ be a torsion-free convex cocompact subgroup of L ; if $\mathbf{k}$ is non-Archimedean, let Γ be any torsion-free finitely generated discrete subgroup of L . Then there is a neighborhood $\mathcal{U} \subset \mathrm{Hom}(\Gamma, G)$ of the natural inclusion such that any $\varphi \in \mathcal{U}$ is injective and $\varphi(\Gamma)$ is discrete in G , acting properly discontinuously on G/H .*

Recall that for $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, a discrete subgroup of L is called *convex cocompact* if it acts cocompactly on the convex hull of its limit set in the

symmetric space of L . In particular, any uniform lattice of L is convex cocompact.

For $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, Theorem 1.1 follows from Theorem 1.3 and from a cohomological argument due to Kobayashi (see Section 5.4). This argument does not transpose to the non-Archimedean case.

Note that in characteristic zero, every finitely generated subgroup of L is virtually torsion-free by Selberg's lemma ([Sel], Lem. 8), hence the "torsion-free" assumption in Theorem 1.3 may easily be removed in this case.

1.3. Translation in terms of a Cartan projection. Let $\mathbf{k}$ be a local field and G the set of $\mathbf{k}$ -points of a connected reductive algebraic $\mathbf{k}$ -group. Fix a Cartan projection $\mu : G \rightarrow E^+$ of G , where E^+ is a closed convex cone in a real finite-dimensional vector space E (see Section 2). For any closed subgroup H of G , the *properness criterion* of Benoist ([Ben], Cor. 5.2) and Kobayashi ([Ko2], Th. 1.1) translates the properness of the action on G/H of a subgroup Γ of G in terms of μ . Using this criterion (see Subsection 5.4), Theorem 1.3 is a consequence of the following result, where we fix a norm $\|\cdot\|$ on E .

Theorem 1.4. *Let $\mathbf{k}$ be a local field, G the set of $\mathbf{k}$ -points of a connected reductive algebraic $\mathbf{k}$ -group $\mathbf{G}$, and L the set of $\mathbf{k}$ -points of a closed reductive subgroup $\mathbf{L}$ of $\mathbf{G}$ of $\mathbf{k}$ -rank 1. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, let Γ be a convex cocompact subgroup of L ; if $\mathbf{k}$ is non-Archimedean, let Γ be any finitely generated discrete subgroup of L . For any $\varepsilon > 0$, there is a neighborhood $\mathcal{U}_\varepsilon \subset \text{Hom}(\Gamma, G)$ of the natural inclusion and a constant $C_\varepsilon \geq 0$ such that*

$$\|\mu(\varphi(\gamma)) - \mu(\gamma)\| \leq \varepsilon \|\mu(\gamma)\| + C_\varepsilon$$

for all $\varphi \in \mathcal{U}_\varepsilon$ and all $\gamma \in \Gamma$.

1.4. Ideas of proofs. The core of the paper is the proof of Theorem 1.4. We start by recalling, in Section 2, that certain linear forms ℓ on E are connected to representations (V, ρ) of $\mathbf{G}$ by relations of the form

$$\ell(\mu(g)) = \log \|\rho(g)\|_V$$

for all $g \in G$, where $\|\cdot\|_V$ is a certain fixed norm on V . We are thus led to bound ratios of the form $\|\rho(\varphi(\gamma))\|_V / \|\rho(\gamma)\|_V$, where $\gamma \in \Gamma \setminus \{1\}$ and where $\varphi \in \text{Hom}(\Gamma, G)$ is close to the natural inclusion of Γ in G .

In order to bound these ratios we look at the dynamics of G acting on the projective space $\mathbb{P}(V)$, notably the dynamics of the elements $g \in G$ that are *proximal* in $\mathbb{P}(V)$. By definition, such elements $g \in G$ admit an attracting fixed point and a repelling projective hyperplane in $\mathbb{P}(V)$. In Section 3 we consider products $z_1 k_2 z_2 \dots k_n z_n$ of proximal elements z_i having a common attracting fixed point x_0^+ and a common repelling hyperplane X_0^- , with isometries k_i such that $k_i \cdot x_0^+$ remains bounded away from X_0^- . We estimate the contraction power of such a product in terms of the contraction powers of the z_i .

In Section 4 we see how such dynamical considerations apply to the elements $\gamma \in \Gamma$ and their images $\varphi(\gamma)$ under a small deformation $\varphi \in \text{Hom}(\Gamma, G)$. We use Guichard's idea [Gui] of writing every element $\gamma \in \Gamma$ as a product $\gamma_0 \dots \gamma_n$ of elements of a fixed finite subset F of Γ , where the norms $\|\mu(\gamma_i)\|$ and $\|\mu(\gamma_i \gamma_{i+1}) - \mu(\gamma_i) - \mu(\gamma_{i+1})\|$ are controlled for all i .

In Section 5 we combine the results of Sections 3 and 4 by carefully choosing the finite subset F of Γ in order to get a sharp control of the ratios $\|\rho(\varphi(\gamma))\|_V / \|\rho(\gamma)\|_V$, or equivalently of $\ell(\mu(\varphi(\gamma)) - \mu(\gamma))$ for $\gamma \in \Gamma \setminus \{1\}$. From this we deduce Theorem 1.4.

At the end of Section 5 we explain how Theorems 1.1 and 1.3 follow from Theorem 1.4. Finally, in Section 6 we establish Corollary 1.2 by relating Theorem 1.1 to Johnson and Millson's bending construction.

Acknowledgements. I warmly thank Yves Benoist and Olivier Guichard for fruitful discussion.

2. CARTAN PROJECTIONS, MAXIMAL PARABOLIC SUBGROUPS, AND REPRESENTATIONS

Throughout the paper, $\mathbf{k}$ denotes a local field, *i.e.*, $\mathbb{R}$, $\mathbb{C}$, a finite extension of $\mathbb{Q}_p$, or the field $\mathbb{F}_q((t))$ of formal Laurent series over a finite field $\mathbb{F}_q$. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, we denote by $|\cdot|$ the usual absolute value on $\mathbf{k}$. If $\mathbf{k}$ is non-Archimedean, we denote by $\mathcal{O}$ the ring of integers of $\mathbf{k}$, by q the cardinal of its residue field, by π a uniformizer, by ω the (additive) valuation on $\mathbf{k}$ such that $\omega(\pi) = 1$, and by $|\cdot| = q^{-\omega(\cdot)}$ the corresponding (multiplicative) absolute value. If $\mathbf{G}$ is an algebraic group, we denote by G the set of its $\mathbf{k}$ -points and by $\mathfrak{g}$ its Lie algebra.

In this section, we recall a few well-known facts on connected reductive algebraic $\mathbf{k}$ -groups and their Cartan projections.

2.1. Weyl chambers. Fix a connected reductive algebraic $\mathbf{k}$ -group $\mathbf{G}$. The derived group $\mathbf{D}(\mathbf{G})$ is semisimple, the identity component $\mathbf{Z}(\mathbf{G})^\circ$ of the center of $\mathbf{G}$ is a torus, which is trivial if $\mathbf{G}$ is semisimple, and $\mathbf{G}$ is the almost product of $\mathbf{D}(\mathbf{G})$ and $\mathbf{Z}(\mathbf{G})^\circ$. Recall that the $\mathbf{k}$ -split $\mathbf{k}$ -tori of $\mathbf{G}$ are all conjugate over $\mathbf{k}$. Fix such a torus $\mathbf{A}$ and let $\mathbf{N}$ (resp. $\mathbf{Z}$) denote its normalizer (resp. centralizer) in $\mathbf{G}$. The group $X(\mathbf{A})$ of $\mathbf{k}$ -characters of $\mathbf{A}$ and the group $Y(\mathbf{A})$ of $\mathbf{k}$ -cocharacters are both free $\mathbb{Z}$ -modules of rank $\text{rank}_{\mathbf{k}}(\mathbf{G})$ and there is a perfect pairing

$$\langle \cdot, \cdot \rangle : X(\mathbf{A}) \times Y(\mathbf{A}) \longrightarrow \mathbb{Z}.$$

Note that $\mathbf{A}$ is the almost product of $(\mathbf{A} \cap \mathbf{D}(\mathbf{G}))^\circ$ and $(\mathbf{A} \cap \mathbf{Z}(\mathbf{G}))^\circ$, hence $X(\mathbf{A}) \otimes_{\mathbb{Z}} \mathbb{R}$ is the direct sum of $X((\mathbf{A} \cap \mathbf{D}(\mathbf{G}))^\circ) \otimes_{\mathbb{Z}} \mathbb{R}$ and $X((\mathbf{A} \cap \mathbf{Z}(\mathbf{G}))^\circ) \otimes_{\mathbb{Z}} \mathbb{R}$.

The set $\Phi = \Phi(\mathbf{A}, \mathbf{G})$ of restricted roots of $\mathbf{A}$ in $\mathbf{G}$, *i.e.*, the set of non-trivial weights of $\mathbf{A}$ in the adjoint representation of $\mathbf{G}$, is a root system of $X((\mathbf{A} \cap \mathbf{D}(\mathbf{G}))^\circ) \otimes_{\mathbb{Z}} \mathbb{R}$. For $\alpha \in \Phi$, let $\check{\alpha}$ be the corresponding coroot: by definition, $\langle \alpha, \check{\alpha} \rangle = 2$ and $s_\alpha(\Phi) = \Phi$, where s_α is the reflection of $X(\mathbf{A}) \otimes_{\mathbb{Z}} \mathbb{R}$ mapping x to $x - \langle x, \check{\alpha} \rangle \alpha$. The group $W = N/Z$ is finite and identifies with the Weyl group of Φ , generated by the reflections s_α .

Similarly, $E = Y(\mathbf{A}) \otimes_{\mathbb{Z}} \mathbb{R}$ is the direct sum of $E_D = Y((\mathbf{A} \cap \mathbf{D}(\mathbf{G}))^\circ) \otimes_{\mathbb{Z}} \mathbb{R}$ and $E_Z = Y((\mathbf{A} \cap \mathbf{Z}(\mathbf{G}))^\circ) \otimes_{\mathbb{Z}} \mathbb{R}$. The group $W = N/Z$ acts trivially on E_Z and identifies with the Weyl group of the root system $\check{\Phi} = \{\check{\alpha}, \alpha \in \Phi\}$ of E_D . We refer to [BoT] for proofs and more detail.

If $\mathbf{k}$ is non-Archimedean, set $A^\circ = A$; if $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, set

$$A^\circ = \{a \in A, \quad \chi(a) \in]0, +\infty[\quad \forall \chi \in X(\mathbf{A})\}.$$

Choose a basis Δ of Φ and let

$$\begin{aligned} A^+ &= \{a \in A^\circ, \quad |\alpha(a)| \geq 1 \quad \forall \alpha \in \Delta\} \\ (\text{resp. } E^+ &= \{x \in E, \quad \langle \alpha, x \rangle \geq 0 \quad \forall \alpha \in \Delta\}) \end{aligned}$$

denote the corresponding closed positive Weyl chamber in A° (resp. in E). The set E^+ is a closed convex cone in the real vector space E . If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, then E identifies with $\mathfrak{a}$ and E^+ with $\log A^+ \subset \mathfrak{a}$, and we endow E with the Euclidean norm $\|\cdot\|$ induced by the Killing form of $\mathfrak{g}$. If $\mathbf{k}$ is non-Archimedean, we endow E with any W -invariant Euclidean norm $\|\cdot\|$.

2.2. Cartan decompositions and Cartan projections. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, there is a maximal compact subgroup K of G such that the Cartan decomposition $G = KA^+K$ holds: for $g \in G$, there are elements $k_g, \ell_g \in K$ and a unique $a_g \in A^+$ such that $g = k_g a_g \ell_g$ ([Hel], Chap. 9, Th. 1.1). Setting $\mu(g) = \log a_g$ defines a map $\mu : G \rightarrow E^+ \simeq \log A^+$, which is continuous, proper, and surjective. It is called the *Cartan projection* with respect to the Cartan decomposition $G = KA^+K$.

If $\mathbf{k}$ is non-Archimedean, let $\text{Res} : X(\mathbf{Z}) \rightarrow X(\mathbf{A})$ denote the restriction homomorphism, where $X(\mathbf{Z})$ is the group of $\mathbf{k}$ -characters of $\mathbf{Z}$. There is a unique group homomorphism $\nu : Z \rightarrow E$ such that

$$\langle \text{Res}(\chi), \nu(z) \rangle = -\omega(\chi(z))$$

for all $\chi \in X(\mathbf{Z})$ and $z \in Z$. Let $Z^+ \subset Z$ denote the inverse image of E^+ under ν . The *Cartan decomposition* $G = KZ^+K$ holds: for $g \in G$, there are elements $k_g, \ell_g \in K$ and $z_g \in Z^+$ such that $g = k_g z_g \ell_g$, and $\nu(z_g)$ is uniquely defined. Setting $\mu(g) = \nu(z_g)$ defines a map $\mu : G \rightarrow E^+$, which is continuous and proper, and whose image $\mu(G)$ is the intersection of E^+ with a lattice of E . It is called the *Cartan projection* with respect to the Cartan decomposition $G = KZ^+K$. For proofs and more detail we refer to the original articles [BT1] and [BT2], but the reader may also find [Rou] a useful reference.

2.3. A geometric interpretation. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, let $X = G/K$ denote the Riemannian symmetric space of G and d its distance; set $x_0 = K \in X$. Since G acts on X by isometries, we have

$$(2.1) \quad \|\mu(g)\| = d(x_0, g \cdot x_0)$$

for all $g \in G$. If $\mathbf{k}$ is non-Archimedean, let X denote the *Bruhat-Tits building* of G : it is a metric space on which G acts properly by isometries with a compact fundamental domain (see [BT1] or [Rou]). When $\text{rank}_{\mathbf{k}}(\mathbf{G}) = 1$, it is a bipartite simplicial tree (see [Ser], § II.1, for the case of $G = \text{SL}_2(\mathbf{k})$). The group K is the stabilizer of some point $x_0 \in X$, and we have

$$(2.2) \quad \|\mu(g)\| = d(x_0, g \cdot x_0)$$

for all $g \in G$, where d denotes the distance on X . It follows from (2.1) and (2.2) that in both cases (Archimedean or not),

$$(2.3) \quad \|\mu(gg')\| \leq \|\mu(g)\| + \|\mu(g')\|$$

for all $g, g' \in G$. In fact, the following stronger inequalities hold (see for instance [Kas], Lem. 2.3): for all $g, g' \in G$,

$$(2.4) \quad \begin{cases} \|\mu(gg') - \mu(g')\| \leq \|\mu(g)\|, \\ \|\mu(gg') - \mu(g)\| \leq \|\mu(g')\|. \end{cases}$$

2.4. Maximal parabolic subgroups. For $\alpha \in \Phi$, let $\mathbf{U}_\alpha$ denote the corresponding unipotent subgroup of $\mathbf{G}$, with Lie algebra $\mathfrak{u}_\alpha = \mathfrak{g}_\alpha \oplus \mathfrak{g}_{2\alpha}$, where

$$\mathfrak{g}_{i\alpha} = \{X \in \mathfrak{g}, \quad \text{Ad}(a)(X) = \alpha(a)^i X \quad \forall a \in A\}$$

for $i = 1, 2$. For any subset θ of Δ , let $\mathbf{P}_\theta$ denote the corresponding *standard* parabolic subgroup of $\mathbf{G}$, with Lie algebra

$$\mathfrak{p}_\theta = \mathfrak{z} \oplus \left(\bigoplus_{\beta \in \Phi^+} \mathfrak{u}_\beta \right) \oplus \left(\bigoplus_{\beta \in \mathbf{N}(\Delta \setminus \theta)} \mathfrak{u}_{-\beta} \right).$$

Every parabolic $\mathbf{k}$ -subgroup $\mathbf{P}$ of $\mathbf{G}$ is conjugate over $\mathbf{k}$ to a unique standard one. In particular, the maximal proper parabolic $\mathbf{k}$ -subgroups of $\mathbf{G}$ are the conjugates of the groups $\mathbf{P}_\alpha = \mathbf{P}_{\{\alpha\}}$, where $\alpha \in \Delta$.

Fix $\alpha \in \Delta$. Since $\mathbf{P}_\alpha$ is its own normalizer in $\mathbf{G}$, the *flag variety* $\mathbf{G}/\mathbf{P}_\alpha$ parametrizes the set of parabolic $\mathbf{k}$ -subgroups that are conjugate to $\mathbf{P}_\alpha$. It is a projective variety, which is defined over $\mathbf{k}$. Let $\mathbf{N}_\alpha^-$ denote the unipotent subgroup of $\mathbf{G}$ generated by the groups $\mathbf{U}_{-\beta}$ for $\beta \in \alpha + \Phi^+$, with Lie algebra

$$\mathfrak{n}_\alpha^- = \bigoplus_{\beta \in \Phi^+} \mathfrak{u}_{-(\alpha+\beta)}.$$

Let W_α be the subgroup of W generated by the reflections s_β for $\beta \in \Delta \setminus \{\alpha\}$. The *Bruhat decomposition*

$$\mathbf{G}/\mathbf{P}_\alpha = \coprod_{w \in W/W_\alpha} \mathbf{N}_\alpha^- w \mathbf{P}_\alpha$$

holds, where the projective subvariety $\mathbf{N}_\alpha^- w \mathbf{P}_\alpha$ has positive codimension whenever $wW_\alpha \neq W_\alpha$. We refer to [BoT] for proofs and more detail.

2.5. Representations of $\mathbf{G}$. For $\alpha \in \Delta$, let $\omega_\alpha \in X(\mathbf{A})$ denote the corresponding fundamental weight: by definition, $\langle \omega_\alpha, \check{\alpha} \rangle = 1$ and $\langle \omega_\alpha, \check{\beta} \rangle = 0$ for all $\beta \in \Delta \setminus \{\alpha\}$. By [Tit], Th. 7.2, there is an irreducible $\mathbf{k}$ -representation (ρ_α, V_α) of $\mathbf{G}$ whose highest weight χ_α is a positive multiple of ω_α and whose highest weight space x_α^+ is a line. The point $x_\alpha^+ \in \mathbb{P}(V_\alpha)$ is the unique fixed point of P_α in $\mathbb{P}(V_\alpha)$. The map from $\mathbf{G}/\mathbf{P}_\alpha$ to $\mathbb{P}(V_\alpha)$ sending $g\mathbf{P}_\alpha$ to $\rho_\alpha(g)(x_\alpha^+)$ is a closed immersion. We denote the set of restricted roots of (ρ_α, V_α) by Λ_α and, for every $\lambda \in \Lambda_\alpha$, the weight space of λ by $(V_\alpha)_\lambda$.

If $\mathbf{k} = \mathbb{R}$ (resp. if $\mathbf{k} = \mathbb{C}$), then the weight spaces are orthogonal with respect to some K -invariant Euclidean (resp. Hermitian) norm $\|\cdot\|_\alpha$ on V_α . The corresponding operator norm $\|\cdot\|_\alpha$ on $\text{End}(V_\alpha)$ satisfies

$$(2.5) \quad \|\rho_\alpha(g)\|_\alpha = e^{\langle \chi_\alpha, \mu(g) \rangle}$$

for all $g \in G$. If $\mathbf{k}$ is non-Archimedean, then there is a K -invariant ultrametric norm $\|\cdot\|_\alpha$ on V_α such that

$$\left\| \sum_{\lambda \in \Lambda_\alpha} v_\lambda \right\|_\alpha = \max_{\lambda \in \Lambda_\alpha} \|v_\lambda\|_\alpha$$

for all $(v_\lambda) \in \prod_{\lambda \in \Lambda_\alpha} (V_\alpha)_\lambda$ and such that the restriction of $\rho_\alpha(z)$ to $(V_\alpha)_\lambda$ is a homothety of ratio $q^{\langle \lambda, \nu(z) \rangle}$ for all $z \in Z$ and all $\lambda \in \Lambda_\alpha$ ([Qui], Th. 6.1). The corresponding operator norm $\|\cdot\|_\alpha$ on $\text{End}(V_\alpha)$ satisfies

$$(2.6) \quad \|\rho_\alpha(g)\|_\alpha = q^{\langle \chi_\alpha, \mu(g) \rangle}$$

for all $g \in G$.

2.6. The example of $\mathbf{SL}_n$. Let $\mathbf{G} = \mathbf{SL}_n$ for some integer $n \geq 2$. The group $\mathbf{A}$ of diagonal matrices with determinant 1 is a maximal $\mathbf{k}$ -split $\mathbf{k}$ -torus of $\mathbf{G}$ which is its own centralizer, *i.e.*, $\mathbf{Z} = \mathbf{A}$. The corresponding root system Φ is the set of linear forms $\varepsilon_i - \varepsilon_j$, $1 \leq i \neq j \leq n$, where

$$\varepsilon_i(\text{diag}(a_1, \dots, a_n)) = a_i.$$

The roots $\varepsilon_i - \varepsilon_{i+1}$, for $1 \leq i \leq n-1$, form a basis Δ of Φ . If $\mathbf{k}$ is Archimedean (resp. non-Archimedean), the corresponding positive Weyl chamber is

$$\begin{aligned} A^+ &= \{ \text{diag}(a_1, \dots, a_n) \in A, \ a_i \in]0, +\infty[\ \forall i \text{ and } a_1 \geq \dots \geq a_n \} \\ (\text{resp. } A^+ &= \{ \text{diag}(a_1, \dots, a_n) \in A, \ |a_1| \geq \dots \geq |a_n| \}). \end{aligned}$$

Set $K = \text{SO}(n)$ (resp. $K = \text{SU}(n)$, resp. $K = \text{SL}_n(\mathcal{O})$) if $\mathbf{k} = \mathbb{R}$ (resp. if $\mathbf{k} = \mathbb{C}$, resp. if $\mathbf{k}$ is non-Archimedean). The Cartan decomposition $G = KA^+K$ holds. If $\mathbf{k} = \mathbb{R}$ (resp. if $\mathbf{k} = \mathbb{C}$) it follows from the polar decomposition in $\text{GL}_n(\mathbb{R})$ (resp. in $\text{GL}_n(\mathbb{C})$) and from the reduction of symmetric (resp. Hermitian) matrices. If $\mathbf{k}$ is non-Archimedean, it follows from the structure theorem for finitely generated modules over a principal ideal domain. The real vector space

$$E = \{ (x_1, \dots, x_n) \in \mathbb{R}^n, \ x_1 + \dots + x_n = 0 \} \simeq \mathbb{R}^{n-1}$$

and its closed convex cone

$$E^+ = \{ (x_1, \dots, x_n) \in E, \ x_1 \geq \dots \geq x_n \}$$

do not depend on $\mathbf{k}$. Let $\mu : G \rightarrow E^+$ denote the Cartan projection with respect to the Cartan decomposition $G = KA^+K$. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, then $\mu(g) = (\frac{1}{2} \log x_i)_{1 \leq i \leq n}$ where x_i is the i -th eigenvalue of ${}^t g g$. If $\mathbf{k}$ is non-Archimedean and if m is any integer such that $\pi^m g \in M_n(\mathcal{O})$, then $\mu(g) = (\omega(x_{m,i}) - m)_{1 \leq i \leq n}$ where $x_{m,i}$ is the i -th invariant factor of $\pi^m g$.

Fix a simple root $\alpha = \varepsilon_{i_0} - \varepsilon_{i_0+1} \in \Delta$. The parabolic group $\mathbf{P}_\alpha$ is defined by the vanishing of the (i, j) -matrix entries for $1 \leq j \leq i_0 < i \leq n$. The flag variety $\mathbf{G}/\mathbf{P}_\alpha$ is the Grassmannian $\mathcal{G}(i_0, n)$ of i_0 -dimensional subspaces of the affine space $\mathbb{A}^n$. The Lie algebra $\mathfrak{n}_\alpha^-$ is defined by the vanishing of the (i, j) -matrix entries for $1 \leq i \leq i_0$ and for $i_0 + 1 \leq i, j \leq n$. The decomposition

$$\mathbf{G}/\mathbf{P}_\alpha = \coprod_{w \in W/W_\alpha} \mathbf{N}_\alpha^- w \mathbf{P}_\alpha$$

is the decomposition of the Grassmannian $\mathcal{G}(i_0, n)$ into Schubert cells. The representation (ρ_α, V_α) is the natural representation of $\mathbf{SL}_n$ in the wedge product $\Lambda^{i_0} \mathbb{A}^n$. Its highest weight is the fundamental weight

$$\omega_\alpha = \varepsilon_1 + \dots + \varepsilon_{i_0}$$

associated with α . The embedding of the Grassmannian $\mathcal{G}(i_0, n)$ into the projective space $\mathbb{P}(V_\alpha) = \mathbb{P}(\Lambda^{i_0} \mathbb{A}^n)$ is the Plücker embedding.

3. DYNAMICS IN PROJECTIVE SPACES

In this section we look at the dynamics of certain endomorphisms of $\mathbf{k}$ -vector spaces in the corresponding projective spaces, where $\mathbf{k}$ is a local field. In Subsection 3.1 we start by recalling the notion of proximality. We then consider products of the form $z_1 k_2 z_2 \dots k_n z_n$, where the z_i are proximal elements with a common attracting fixed point x_0^+ and a common repelling hyperplane X_0^- , and the k_i are isometries such that $k_i \cdot x_0^+$ remains bounded away from X_0^- . We estimate the contraction power of such a product in terms of the contraction powers of the z_i . In Subsection 3.2 we consider a connected reductive algebraic $\mathbf{k}$ -group $\mathbf{G}$ and apply the result of Subsection 3.1 to the representations (V_α, ρ_α) of $\mathbf{G}$ introduced in Subsection 2.5. From (2.5) and (2.6) we get an upper bound for $|\langle \chi_\alpha, \mu(g_1 \dots g_n) - \mu(g_1) - \dots - \mu(g_n) \rangle|$ for elements $g_1, \dots, g_n \in G$ satisfying certain contractivity and transversality conditions.

3.1. Proximality in projective spaces and norm estimates. Let $\mathbf{k}$ be a local field and V be a finite-dimensional vector space over $\mathbf{k}$. Given a basis $(v_1, \dots, v_n)$ of V , we define the norm

$$(3.1) \quad \left\| \sum_{1 \leq j \leq n} t_j v_j \right\|_V = \sup_{1 \leq j \leq n} |t_j|$$

on V , and we keep the notation $\|\cdot\|_V$ for the corresponding operator norm on $\text{End}(V)$. We endow the projective space $\mathbb{P}(V)$ with the distance

$$d(x_1, x_2) = \inf \{ \|v_1 - v_2\|_V, v_i \in x_i \text{ and } \|v_i\|_V = 1 \ \forall i = 1, 2 \}.$$

Recall that an element $g \in \text{End}(V) \setminus \{0\}$ is called *proximal* if it has a unique eigenvalue of maximal absolute value and if this eigenvalue has multiplicity 1. (The eigenvalues of g belong to a finite extension $\mathbf{k}_g$ of $\mathbf{k}$ and we consider the unique extension to $\mathbf{k}_g$ of the absolute value $|\cdot|$ on $\mathbf{k}$.) If g is proximal, then its maximal eigenvalue belongs to $\mathbf{k}$; we denote by $x_g^+ \in \mathbb{P}(V)$ the corresponding eigenline and by X_g^- the image in $\mathbb{P}(V)$ of the unique g -invariant complementary subspace of x_g^+ in V . Note that g acts on $\mathbb{P}(V)$ by contracting $\mathbb{P}(V) \setminus X_g^-$ towards x_g^+ . For $\varepsilon > 0$, we will say that g is ε -proximal if it satisfies the two following additional conditions:

- (1) $d(x_g^+, X_g^-) \geq 2\varepsilon$,
- (2) for any $x \in \mathbb{P}(V)$, if $d(x, X_g^-) \geq \varepsilon$, then $d(g \cdot x, x_g^+) \leq \varepsilon$.

We will need the following lemma.

Lemma 3.1. *Let X_0^- be a projective hyperplane of $\mathbb{P}(V)$, let $x_0^+ \in \mathbb{P}(V) \setminus X_0^-$, and let $\varepsilon > 0$ such that $d(x_0^+, X_0^-) \geq 2\varepsilon$. Then there exists $r_\varepsilon > 0$ such that for any isometries $k_2, \dots, k_n \in \text{End}(V)$ with $d(k_i \cdot x_0^+, X_0^-) \geq 2\varepsilon$ and for any ε -proximal endomorphisms $z_1, \dots, z_n \in \text{End}(V)$ with $x_{z_i}^+ = x_0^+$ and $X_{z_i}^- = X_0^-$, inducing a homothety of ratio $\|z_i\|_V$ on the line x_0^+ , we have*

$$e^{-(n-1)r_\varepsilon} \cdot \prod_{i=1}^n \|z_i\|_V \leq \|z_1 k_2 z_2 \dots k_n z_n\|_V \leq \prod_{i=1}^n \|z_i\|_V.$$

Proof. Since the operator norm $\|\cdot\|_V$ on $\text{End}(V)$ is submultiplicative and k_i is an isometry of V for all i ,

$$\|z_1 k_2 z_2 \dots k_n z_n\|_V \leq \prod_{i=1}^n \|z_i\|_V.$$

Let us prove the left-hand inequality. Let $v_0 \in V \setminus \{0\}$ satisfy $x_0^+ = \mathbf{k}v_0$ and let V_0 be the hyperplane of V such that $X_0^- = \mathbb{P}(V_0)$. Set

$$\begin{aligned} b_\varepsilon &= \{x \in \mathbb{P}(V), d(x, x_0^+) \leq \varepsilon\} \\ \text{and } B_\varepsilon &= \{x \in \mathbb{P}(V), d(x, X_0^-) \geq \varepsilon\}. \end{aligned}$$

Note that the set of unitary vectors $v \in V$ with $\mathbf{k}v \in B_\varepsilon$ is compact and that the map sending $v \in V$ to $t \in \mathbf{k}$ such that $v \in tv_0 + V_0$ is continuous, hence there exists $r_\varepsilon > 0$ such that

$$(3.2) \quad v \in [e^{-\frac{r_\varepsilon}{2}}, e^{\frac{r_\varepsilon}{2}}] \|v\|_V v_0 + V_0$$

for all $v \in V \setminus \{0\}$ with $\mathbf{k}v \in B_\varepsilon$. Set

$$h_j = z_j k_{j+1} z_{j+1} \dots k_n z_n \in \text{End}(V)$$

for $1 \leq j \leq n$. We claim that $h_j \cdot B_\varepsilon \subset b_\varepsilon$ and

$$(3.3) \quad \|h_j \cdot v_0\|_V \geq e^{-(n-j)r_\varepsilon} \cdot \prod_{i=j}^n \|z_i\|_V$$

for all j . This follows from an easy descending induction on j . Indeed, for all i we have $k_i \cdot b_\varepsilon \subset B_\varepsilon$ since k_i is an isometry of V and $d(k_i \cdot x_0^+, X_0^-) \geq 2\varepsilon$, and $z_i \cdot B_\varepsilon \subset b_\varepsilon$ since z_i is ε -proximal with $x_{z_i}^+ = x_0^+$ and $X_{z_i}^- = X_0^-$. By (3.2), we have $k_{j+1} h_{j+1} \cdot v_0 \in t_j v_0 + V_0$ for some $t_j \in \mathbb{R}$ with

$$|t_j| \geq e^{-\frac{r_\varepsilon}{2}} \|k_{j+1} h_{j+1} \cdot v_0\|_V = e^{-\frac{r_\varepsilon}{2}} \|h_{j+1} \cdot v_0\|_V.$$

By the inductive assumption,

$$|t_j| \geq e^{-(n-j-\frac{1}{2})r_\varepsilon} \cdot \prod_{i=j+1}^n \|z_i\|_V.$$

By hypothesis, z_j preserves V_0 and induces a homothety of ratio $\|z_j\|_V$ on the line x_0^+ , hence $h_j \cdot v_0 = z_j k_{j+1} h_{j+1} \cdot v_0 \in \|z_j\|_V t_j v_0 + V_0$, where

$$\|z_j\|_V |t_j| \geq e^{-(n-j-\frac{1}{2})r_\varepsilon} \cdot \prod_{i=j}^n \|z_i\|_V.$$

Inequality (3.3) follows, using (3.2) again. $\square$

3.2. Cartan projection along the fundamental weights. Lemma 3.1 implies the following result.

Proposition 3.2. *Let $\mathbf{k}$ be a local field and G the set of $\mathbf{k}$ -points of a connected reductive algebraic $\mathbf{k}$ -group. Let $G = KA^+K$ or $G = KZ^+K$ be a Cartan decomposition and $\mu : G \rightarrow E^+$ the corresponding Cartan projection. Fix $\alpha \in \Delta$ and let $\mathcal{C}_\alpha$ be a compact subset of N_α^- . Then there exist $r_\alpha, R_\alpha > 0$*

such that for all $g_1, \dots, g_n \in G$ with $\langle \alpha, \mu(g_i) \rangle \geq R_\alpha$ and $\ell_{g_i} k_{g_{i+1}} \in \mathcal{C}_\alpha P_\alpha$, we have

$$\left| \left\langle \chi_\alpha, \mu(g_1 \dots g_n) - \sum_{i=1}^n \mu(g_i) \right\rangle \right| \leq nr_\alpha.$$

We keep notation from Section 2. In particular, for $g \in G$ we write $g = k_g z_g \ell_g$ with $k_g, \ell_g \in K$ and $z_g \in Z^+$, as in Subsection 2.2. Given a simple root $\alpha \in \Delta$, we denote by $\mathbf{N}_\alpha^-$ (resp. by $\mathbf{P}_\alpha$) the unipotent (resp. parabolic) subgroup of $\mathbf{G}$ introduced in Subsection 2.4, and by χ_α the highest weight of the representation (V_α, ρ_α) introduced in Subsection 2.5.

Precisely, Proposition 3.2 follows from Lemma 3.1, from (2.5) and (2.6), and from the following lemma.

Lemma 3.3. *Let $x_\alpha^+ \in \mathbb{P}(V_\alpha)$ be the highest weight line $(V_\alpha)_{\chi_\alpha}$, and let X_α^- be the image in $\mathbb{P}(V_\alpha)$ of the sum of the weight spaces $(V_\alpha)_\lambda$ for $\lambda \in \Lambda_\alpha \setminus \{\chi_\alpha\}$.*

- (1) *Given $\varepsilon > 0$ with $d(x_\alpha^+, X_\alpha^-) \geq 2\varepsilon$, there exists $R_\alpha > 0$ such that for any $z \in Z^+$ with $\langle \alpha, \mu(z) \rangle \geq R_\alpha$, the element $\rho_\alpha(z)$ is ε -proximal in $\mathbb{P}(V_\alpha)$ with $(x_{\rho_\alpha(z)}^+, X_{\rho_\alpha(z)}^-) = (x_\alpha^+, X_\alpha^-)$.*
- (2) *We have $\rho_\alpha(N_\alpha^-)(x_\alpha^+) \cap X_\alpha^- = \emptyset$.*

Proof. (1) It is sufficient to see that every restricted weight of (ρ_α, V_α) except χ_α belongs to $\chi_\alpha - \alpha - \mathbb{N}\Delta$. Consider the subgroup W_α of W generated by the reflections $s_\beta : x \mapsto x - \langle x, \check{\beta} \rangle \beta$ for $\beta \in \Delta \setminus \{\alpha\}$. It acts transitively on the root subsystem of Φ generated by $\Delta \setminus \{\alpha\}$, and it fixes χ_α since χ_α is a multiple of ω_α and $\langle \omega_\alpha, \check{\beta} \rangle = 0$ for all $\beta \in \Delta \setminus \{\alpha\}$. Therefore, for every weight $\lambda \in \chi_\alpha - \mathbb{N}(\Delta \setminus \{\alpha\})$ there exists $w \in W_\alpha$ such that $w \cdot \lambda \in \chi_\alpha + \mathbb{N}\Delta$, which implies that $\lambda = \chi_\alpha$.

(2) For $n \in N_\alpha^-$, the identity element $1 \in G$ belongs to the closure of the conjugacy class $\{znz^{-1}, z \in Z\}$, hence x_α^+ belongs to the closure of the orbit $\rho_\alpha(Zn)(x_\alpha^+)$ in $\mathbb{P}(V_\alpha)$. But X_α^- is closed in $\mathbb{P}(V_\alpha)$, stable under Z , and does not contain x_α^+ . $\square$

Proof of Proposition 3.2. The point $x_\alpha^+ \in \mathbb{P}(V_\alpha)$ is fixed by P_α . Moreover, $\rho_\alpha(N_\alpha^-)(x_\alpha^+) \cap X_\alpha^- = \emptyset$ by Lemma 3.3, hence there exists $\varepsilon > 0$ such that

$$d(\rho_\alpha(\mathcal{C}_\alpha P_\alpha)(x_\alpha^+), X_\alpha^-) \geq 2\varepsilon.$$

Let R_α be given by Lemma 3.3 and let $r_\alpha = r_\varepsilon / \log q$, where $r_\varepsilon > 0$ is given by Lemma 3.1 and $q = e$ if $\mathbf{k}$ is Archimedean, q is the cardinal of the residue field of $\mathcal{O}$ otherwise. Let $g_1, \dots, g_n \in G$ satisfy $\langle \alpha, \mu(g_i) \rangle \geq R_\alpha$ and $\ell_{g_i} k_{g_{i+1}} \in \mathcal{C}_\alpha P_\alpha$ for all i . By Lemma 3.3, $\rho_\alpha(z_{g_i})$ is ε -proximal in $\mathbb{P}(V_\alpha)$ with $x_{\rho_\alpha(z_{g_i})}^+ = x_\alpha^+$ and $X_{\rho_\alpha(z_{g_i})}^- = X_\alpha^-$. Moreover, it induces a homothety of ratio $\|\rho_\alpha(z_{g_i})\|_\alpha$ on the line x_α^+ . By Lemma 3.1,

$$q^{-nr_\alpha} \cdot \prod_{i=1}^n \|\rho_\alpha(z_{g_i})\|_\alpha \leq \|\rho_\alpha(g_1 \dots g_n)\|_\alpha \leq \prod_{i=1}^n \|\rho_\alpha(z_{g_i})\|_\alpha.$$

Using (2.5) and (2.6), we get

$$\left\langle \chi_\alpha, \sum_{i=1}^n \mu(g_i) \right\rangle - nr_\alpha \leq \langle \chi_\alpha, \mu(g_1 \dots g_n) \rangle \leq \left\langle \chi_\alpha, \sum_{i=1}^n \mu(g_i) \right\rangle. \quad \square$$

4. TRANSVERSE PRODUCTS

In this section we explain how, under the assumptions of Theorem 1.4, Proposition 3.2 applies to the elements $\gamma \in \Gamma$ and their images $\varphi(\gamma)$ under a small deformation $\varphi \in \text{Hom}(\Gamma, G)$. We use Guichard's idea [Gui] of writing every element $\gamma \in \Gamma$ as a “transverse product” $\gamma_0 \dots \gamma_n$ of elements of a fixed finite subset F of Γ .

4.1. Transversality in L . Let $\mathbf{k}$ be a local field and $\mathbf{L}$ a connected reductive algebraic $\mathbf{k}$ -group of $\mathbf{k}$ -rank 1. Fix a Cartan decomposition $L = K_L A_L^+ K_L$ or $L = K_L Z_L^+ K_L$, where K_L is a maximal compact subgroup of L , where $\mathbf{A}_{\mathbf{L}}$ is a maximal $\mathbf{k}$ -split $\mathbf{k}$ -torus of $\mathbf{L}$, and where $\mathbf{Z}_{\mathbf{L}}$ is the centralizer of $\mathbf{A}_{\mathbf{L}}$ in $\mathbf{L}$. Let $\mu_L : L \rightarrow E_L^+$ denote the corresponding Cartan projection, where $E_L = Y(\mathbf{A}_{\mathbf{L}}) \otimes_{\mathbb{Z}} \mathbb{R}$. Since $\mathbf{L}$ has $\mathbf{k}$ -rank 1, the vector space E_L is a line, and any isomorphism from E_L to $\mathbb{R}$ gives a Cartan projection $\mu_L^{\mathbb{R}} : L \rightarrow \mathbb{R}$.

If $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 1, then $\mu_L^{\mathbb{R}}$ takes only nonnegative or only nonpositive values. We denote by α_L the indivisible positive restricted root of $\mathbf{A}_{\mathbf{L}}$ in $\mathbf{L}$, by $\mathbf{P}_{\mathbf{L}} = \mathbf{P}_{\alpha_L}$ the proper parabolic subgroup of $\mathbf{L}$ associated with α_L , and by $\mathbf{N}_{\mathbf{L}}^- = \mathbf{U}_{-\alpha_L}$ the unipotent subgroup associated with $-\alpha_L$.

If $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 0, then $\mathbf{A}_{\mathbf{L}}$ is central in $\mathbf{L}$, hence $\mathbf{Z}_{\mathbf{L}} = \mathbf{L}$. In this case $\mu_L^{\mathbb{R}}$ is a group homomorphism from L to $\mathbb{R}$, thus taking both positive and negative values. We set $\mathbf{P}_{\mathbf{L}} = \mathbf{Z}_{\mathbf{L}} = \mathbf{L}$ and $\mathbf{N}_{\mathbf{L}}^- = \{1\}$.

For the reader's convenience, we give a proof of the following result, which is due to Guichard in the real semisimple case ([Gui], Lem. 7 & 9). We consider the more general situation of a reductive algebraic group over a local field.

Proposition 4.1 (Guichard). *Let $\mathbf{k}$ be a local field, L the set of $\mathbf{k}$ -points of a connected reductive algebraic $\mathbf{k}$ -group of $\mathbf{k}$ -rank 1, and $\mu_L^{\mathbb{R}} : L \rightarrow \mathbb{R}$ a Cartan projection. If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, let Γ be a convex cocompact subgroup of L ; if $\mathbf{k}$ is non-Archimedean, let Γ be any finitely generated discrete subgroup of L . Then there exist $D > 0$ and a compact subset $\mathcal{C}_L$ of N_L^- such that for $R \geq D$, any $\gamma \in \Gamma$ may be written as $\gamma = \gamma_0 \dots \gamma_n$, where*

- (1) $|\mu_L^{\mathbb{R}}(\gamma_0)| \leq R + D$ and $R - D \leq |\mu_L^{\mathbb{R}}(\gamma_i)| \leq R + D$ for all $1 \leq i \leq n$,
- (2) $\mu_L^{\mathbb{R}}(\gamma_1), \dots, \mu_L^{\mathbb{R}}(\gamma_n)$ are all ≥ 0 or all ≤ 0 ,
- (3) $\ell_{\gamma_i} k_{\gamma_{i+1}} \in \mathcal{C}_L P_L$ for all $1 \leq i \leq n - 1$.

To prove Proposition 4.1 we use the following lemma, which translates the transversality condition (3) in terms of $\mu_L^{\mathbb{R}}$.

Lemma 4.2. *Under the assumptions of Proposition 4.1, there exists $D_0 \geq 0$ with the following property: given any $D \geq D_0$, there is a compact subset $\mathcal{C}_L$ of N_L^- such that for $k \in K_L$, if*

$$|\mu_L^{\mathbb{R}}(z_1 k z_2)| \geq |\mu_L^{\mathbb{R}}(z_1)| + |\mu_L^{\mathbb{R}}(z_2)| - D$$

for some $z_1, z_2 \in Z_L^+$ with $|\mu_L^{\mathbb{R}}(z_1)|, |\mu_L^{\mathbb{R}}(z_2)| \geq D$, then $k \in \mathcal{C}_L P_L$.

Note that Proposition 3.2 implies some kind of converse to Lemma 4.2: for any compact subset $\mathcal{C}_L$ of N_L^- , there exists $D \geq 0$ such that for all $k \in K_L \cap \mathcal{C}_L P_L$ and all $z_1, z_2 \in Z_L^+$,

$$|\mu_L^{\mathbb{R}}(z_1 k z_2)| \geq |\mu_L^{\mathbb{R}}(z_1)| + |\mu_L^{\mathbb{R}}(z_2)| - D.$$

Proof of Lemma 4.2. We may assume that $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 1. Then L/P_L is the disjoint union of $N_L^- \cdot P_L$ and $\{w \cdot P_L\}$, where w denotes the nontrivial element of the (restricted) Weyl group of L . It is therefore sufficient to prove the existence of a neighborhood $\mathcal{U}$ of $w \cdot P_L$ in L/P_L such that for all $k \in K_L$ with $k \cdot P_L \in \mathcal{U}$ and all $z_1, z_2 \in Z_L^+$ with $\mu_L^{\mathbb{R}}(z_1), \mu_L^{\mathbb{R}}(z_2) \geq D$, we have

$$|\mu_L^{\mathbb{R}}(z_1 k z_2)| < |\mu_L^{\mathbb{R}}(z_1)| + |\mu_L^{\mathbb{R}}(z_2)| - D.$$

Let X_L denote either the Riemannian symmetric space or the Bruhat-Tits tree of L , depending on whether $\mathbf{k}$ is Archimedean or not. The space X_L is Gromov-hyperbolic and we may identify L/P_L with the boundary at infinity ∂X_L of X_L , *i.e.*, with the set of equivalence classes $[\mathcal{R}]$ of geodesic half-lines $\mathcal{R} : [0, +\infty[\rightarrow X_L$ for the equivalence relation “to stay at bounded distance”. The point $P_L \in L/P_L$ (resp. $w \cdot P_L \in L/P_L$) corresponds to the equivalence class $[\mathcal{R}^+]$ (resp. $[\mathcal{R}^-]$) of the geodesic half-line $\mathcal{R}^+ : [0, +\infty[\rightarrow X$ (resp. $\mathcal{R}^- : [0, +\infty[\rightarrow X$) whose image is $Z_L^+ \cdot x_0$ (resp. $(w \cdot Z_L^+) \cdot x_0$). Let d be the distance on X_L and x_0 the point of X_L whose stabilizer is K_L . By (2.1) and (2.2), we may assume that $|\mu_L^{\mathbb{R}}(g)| = d(x_0, g \cdot x_0)$ for all $g \in L$. By the “shadow lemma” (see [Bou], Lem. 1.6.2, for instance), there is a constant $D_0 > 0$ such that the open sets

$$\mathcal{U}_t = \left\{ [\mathcal{R}], \quad \mathcal{R}(0) = x_0 \text{ and } d(\mathcal{R}(t), \mathcal{R}^-(t)) < D_0 \right\},$$

for $t \in [0, +\infty[$, form a basis of neighborhoods of $[\mathcal{R}^-]$ in ∂X_L . Fix $D \geq D_0$. For all $k \in K_L$ and $z_1, z_2 \in Z_L^+$ with $t_1 := \mu_L^{\mathbb{R}}(z_1) \geq D$ and $t_2 := \mu_L^{\mathbb{R}}(z_2) \geq D$, we have

$$\begin{aligned} |\mu_L^{\mathbb{R}}(z_1 k z_2)| &= d(x_0, z_1 k z_2 \cdot x_0) \\ &= d(z_1^{-1} \cdot x_0, k z_2 \cdot x_0) \\ &= d(\mathcal{R}^-(t_1), k \cdot \mathcal{R}^+(t_2)) \\ &\leq d(\mathcal{R}^-(t_1), \mathcal{R}^-(D)) + d(\mathcal{R}^-(D), k \cdot \mathcal{R}^+(D)) \\ &\quad + d(k \cdot \mathcal{R}^+(D), k \cdot \mathcal{R}^+(t_2)) \\ &= t_1 - D + d(\mathcal{R}^-(D), k \cdot \mathcal{R}^+(D)) + t_2 - D \\ &= |\mu_L^{\mathbb{R}}(z_1)| + |\mu_L^{\mathbb{R}}(z_2)| - 2D + d(\mathcal{R}^-(D), k \cdot \mathcal{R}^+(D)). \end{aligned}$$

Therefore, if $[k \cdot \mathcal{R}^+] \in \mathcal{U}_D$ then $|\mu_L^{\mathbb{R}}(z_1 k z_2)| < |\mu_L^{\mathbb{R}}(z_1)| + |\mu_L^{\mathbb{R}}(z_2)| - D$. This completes the proof of Lemma 4.2. $\square$

Proof of Proposition 4.1. As in the proof of Lemma 4.2, let X_L denote either the Riemannian symmetric space or the Bruhat-Tits tree of L , depending on whether $\mathbf{k}$ is Archimedean or not. Let d be the distance on X_L and x_0 the point of X_L whose stabilizer is K_L . By (2.1) and (2.2), we may assume that $|\mu_L^{\mathbb{R}}(g)| = d(g \cdot x_0, x_0)$ for all $g \in L$. Let X'_L denote the convex hull of the limit set of Γ in X_L . It is a closed subset of X_L on which Γ acts cocompactly: indeed, if $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$ this is the convex cocompactness assumption; if $\mathbf{k}$ is non-Archimedean it follows from [Bas], Prop. 7.9. Fix a compact fundamental domain $\mathcal{D}$ of X'_L for the action of Γ , and fix $x'_0 \in \text{Int}(\mathcal{D})$. Let $d_{\mathcal{D}}$ be the diameter of $\mathcal{D}$ and D_0 the constant given by Lemma 4.2. Let

$$D = \max(D_0, 6 d_{\mathcal{D}} + 6 d(x_0, x'_0)) > 0$$

and let $\mathcal{C}_L$ be the corresponding compact subset of N_L^- given by Lemma 4.2. We claim that D and $\mathcal{C}_L$ satisfy the conclusions of Proposition 4.1. Indeed, let $R \geq D$. Fix $\gamma \in \Gamma$ and let I be the geodesic segment of X'_L with endpoints x'_0 and $\gamma^{-1} \cdot x'_0$. Let $n \in \mathbb{N}$ such that

$$nR \leq d(x'_0, \gamma^{-1} \cdot x'_0) < (n+1)R.$$

For all $1 \leq i \leq n$, let $x'_i \in I$ satisfy $d(x'_i, x'_0) = iR$. We have $x'_i \in \lambda_i \cdot \mathcal{D}$ for some $\lambda_i \in \Gamma$. Let $\gamma_0 = \gamma \lambda_n \in \Gamma$ and $\gamma_i = \lambda_{n-i+1}^{-1} \lambda_{n-i} \in \Gamma$ for $i \geq 1$ (where $\lambda_0 = 1$), so that $\gamma = \gamma_0 \dots \gamma_n$. For all $1 \leq i \leq n$,

$$\begin{aligned} \left| |\mu_L^{\mathbb{R}}(\gamma_i)| - d(x'_{n-i}, x'_{n-i+1}) \right| &= \left| d(\lambda_{n-i} \cdot x_0, \lambda_{n-i+1} \cdot x_0) - d(x'_{n-i}, x'_{n-i+1}) \right| \\ &\leq d(\lambda_{n-i} \cdot x_0, \lambda_{n-i} \cdot x'_0) + d(\lambda_{n-i} \cdot x'_0, x'_{n-i}) \\ &\quad + d(x'_{n-i+1}, \lambda_{n-i+1} \cdot x'_0) + d(\lambda_{n-i+1} \cdot x'_0, \lambda_{n-i+1} \cdot x_0) \\ &\leq 2d_{\mathcal{D}} + 2d(x_0, x'_0). \end{aligned}$$

Since $d(x'_{n-i}, x'_{n-i+1}) = R$, we have $||\mu_L^{\mathbb{R}}(\gamma_i)| - R| \leq 2d_{\mathcal{D}} + 2d(x_0, x'_0)$. Similarly,

$$||\mu_L^{\mathbb{R}}(\gamma_0)| - d(x'_n, \gamma^{-1} \cdot x'_0)| \leq 2d_{\mathcal{D}} + 2d(x_0, x'_0),$$

hence $|\mu_L^{\mathbb{R}}(\gamma_0)| \leq R + 2d_{\mathcal{D}} + 2d(x_0, x'_0)$. For $1 \leq i \leq n-1$, the same reasoning shows that

$$\begin{aligned} |\mu_L^{\mathbb{R}}(\gamma_i \gamma_{i+1})| &\geq d(x_{n-i-1}, x_{n-i+1}) - 2d_{\mathcal{D}} - 2d(x_0, x'_0) \\ &= 2R - 2d_{\mathcal{D}} - 2d(x_0, x'_0) \\ &\geq |\mu_L^{\mathbb{R}}(\gamma_i)| + |\mu_L^{\mathbb{R}}(\gamma_{i+1})| - 6d_{\mathcal{D}} - 6d(x_0, x'_0) \\ (4.1) \quad &\geq |\mu_L^{\mathbb{R}}(\gamma_i)| + |\mu_L^{\mathbb{R}}(\gamma_{i+1})| - D. \end{aligned}$$

By Lemma 4.2, we have $\ell_{\gamma_i} k_{\gamma_{i+1}} \in \mathcal{C}_L P_L$ for all $1 \leq i \leq n-1$.

We claim that $\mu_L^{\mathbb{R}}(\gamma_1), \dots, \mu_L^{\mathbb{R}}(\gamma_n) \in \mathbb{R}$ all have the same sign. Indeed, we may assume that $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 0, in which case $\mu_L^{\mathbb{R}} : L \rightarrow \mathbb{R}$ is a group homomorphism. If $\mu_L^{\mathbb{R}}(\gamma_i)$ and $\mu_L^{\mathbb{R}}(\gamma_{i+1})$ had different signs for some $1 \leq i \leq n-1$, then (4.1) would imply that

$$\min(|\mu_L^{\mathbb{R}}(\gamma_i)|, |\mu_L^{\mathbb{R}}(\gamma_{i+1})|) \leq \frac{D}{2},$$

which would contradict the fact that

$$\begin{aligned} |\mu_L^{\mathbb{R}}(\gamma_i)|, |\mu_L^{\mathbb{R}}(\gamma_{i+1})| &\geq R - 2d_{\mathcal{D}} - 2d(x_0, x'_0) \\ &\geq D - 2d_{\mathcal{D}} - 2d(x_0, x'_0) > \frac{D}{2}. \quad \square \end{aligned}$$

4.2. Transversality in G . Let $\mathbf{k}$ be a local field, $\mathbf{G}$ a connected reductive algebraic $\mathbf{k}$ -group, and $\mathbf{L}$ a closed connected reductive subgroup of $\mathbf{G}$ of $\mathbf{k}$ -rank 1. Fix a Cartan decomposition $G = KA^+K$ or $G = KZ^+K$.

Remark 4.3. After conjugating $\mathbf{L}$ by some element of G , we may assume that L admits a Cartan decomposition $L = K_L A_L^+ K_L$ or $L = K_L Z_L^+ K_L$ with $K_L \subset K$, with $\mathbf{A}_L \subset \mathbf{A}$, and with $A_L^+ \cap A^+$ noncompact.

Indeed, $\mathbf{A}_L$ is contained in some maximal $\mathbf{k}$ -split $\mathbf{k}$ -torus of $\mathbf{G}$, and these tori are all conjugate over $\mathbf{k}$ ([BoT], Th. 4.21). Thus, after conjugating $\mathbf{L}$ by some element of G , we may assume that $\mathbf{A}_L \subset \mathbf{A}$. We now use a result proved by Mostow [Mos] and Karpelevich [Kar] in the Archimedean case,

and by Landvogt [Lan] in the non-Archimedean case: after conjugating $\mathbf{L}$ again by some element of G , we may assume that $K_L \subset K$. Finally, after conjugating $\mathbf{L}$ by some element of the Weyl group W , we may assume that $A_L^+ \cap A^+$ is noncompact.

Assume that the conditions of Remark 4.3 are satisfied. The following lemma provides a link between Propositions 3.2 and 4.1. We use the notation of Subsection 2.4.

Lemma 4.4. *If the restriction of $\alpha \in \Delta$ to $\mathbf{A}_L$ is nontrivial, then $P_L \subset P_\alpha$ and $N_L^- \subset N_\alpha^- P_\alpha$.*

Proof. Fix $\alpha \in \Delta$ whose restriction to $\mathbf{A}_L$ is nontrivial, and let $a \in A_L^+ \cap A^+$ such that $|\alpha(a)| > 1$. Note that $\mathfrak{g} = \mathfrak{n}_\alpha^- \oplus \mathfrak{p}_\alpha$ and $\mathfrak{p}_\alpha = \mathfrak{p}_\emptyset \oplus \mathfrak{n}_{\alpha^c}^-$, where

$$\begin{aligned} \mathfrak{n}_\alpha^- &= \bigoplus_{\beta \in \Phi^+} \mathfrak{u}_{-(\alpha+\beta)}, \\ \mathfrak{p}_\emptyset &= \mathfrak{z} \oplus \bigoplus_{\beta \in \Phi^+} \mathfrak{u}_\beta, \\ \text{and } \mathfrak{n}_{\alpha^c}^- &= \bigoplus_{\beta \in \mathbb{N}(\Delta \setminus \{\alpha\})} \mathfrak{u}_{-\beta} \end{aligned}$$

are all direct sums of eigenspaces of $\text{Ad}(a)$, with eigenvalues of absolute value < 1 on $\mathfrak{n}_\alpha^-$ and ≥ 1 on $\mathfrak{p}_\emptyset$. Since $\mathfrak{p}_L$ is a sum of eigenspaces of $\text{Ad}(a)$ for eigenvalues of absolute value ≥ 1 , we have $\mathfrak{p}_L \subset \mathfrak{p}_\alpha$. Given that $\mathbf{P}_L$ and $\mathbf{P}_\alpha$ are connected, this implies that $P_L \subset P_\alpha$. Since $\mathfrak{n}_L^-$ is a sum of eigenspaces of $\text{Ad}(a)$ for eigenvalues of absolute value < 1 , we have $\mathfrak{n}_L^- \subset \mathfrak{n}_\alpha^- \oplus \mathfrak{n}_{\alpha^c}^-$. Note that $[\mathfrak{n}_\alpha^-, \mathfrak{n}_{\alpha^c}^-] \subset \mathfrak{n}_\alpha^-$, hence N_α^- is normalized by the group $N_{\alpha^c}^-$ generated by the groups $U_{-\beta}$ for $\beta \in \mathbb{N}(\Delta \setminus \{\alpha\})$. This implies that

$$N_L^- \subset N_\alpha^- N_{\alpha^c}^- \subset N_\alpha^- P_\alpha. \quad \square$$

5. CARTAN PROJECTION AND DEFORMATION

In this section we prove Theorem 1.4 using Propositions 3.2 and 4.1. By the triangular inequality, it is sufficient to prove the following proposition.

Proposition 5.1. *Let $\mathbf{k}$ be a local field, G the set of $\mathbf{k}$ -points of a connected reductive algebraic $\mathbf{k}$ -group $\mathbf{G}$, and L the set of $\mathbf{k}$ -points of a closed reductive subgroup $\mathbf{L}$ of $\mathbf{G}$ of $\mathbf{k}$ -rank 1. Fix a Cartan projection $\mu : G \rightarrow E^+$ and a norm $\|\cdot\|$ on E . If $\mathbf{k} = \mathbb{R}$ or $\mathbb{C}$, let Γ be a convex cocompact subgroup of L ; if $\mathbf{k}$ is non-Archimedean, let Γ be any finitely generated discrete subgroup of L . Then for any $\varepsilon > 0$, there exist a finite subset F_ε of Γ , a neighborhood $\mathcal{U}_\varepsilon \subset \text{Hom}(\Gamma, G)$ of the natural inclusion, and a constant $C_\varepsilon \geq 0$ such that any $\gamma \in \Gamma$ may be written as $\gamma = \gamma_0 \dots \gamma_n$ for some $\gamma_0, \dots, \gamma_n \in F_\varepsilon$ with*

- (1) $n \leq \varepsilon \|\mu(\gamma)\| + C_\varepsilon$,
- (2) $\|\mu(\varphi(\gamma_i)) - \mu(\gamma_i)\| \leq 1$ for all $\varphi \in \mathcal{U}_\varepsilon$ and $0 \leq i \leq n$,
- (3) for all $\varphi \in \mathcal{U}_\varepsilon$,

$$\left\| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right\| \leq \varepsilon \|\mu(\gamma)\| + C_\varepsilon.$$

The proof of Proposition 5.1 will be given in Subsection 5.3.

5.1. Norms on E and its subspaces. Under the assumptions of Proposition 5.1, let $G = KA^+K$ or $G = KZ^+K$ be the Cartan decomposition corresponding to μ . By (2.4), in order to prove Proposition 5.1, we may assume that $\mathbf{L}$ is connected and replace it by any conjugate by G . By Remark 4.3, we may assume that L admits a Cartan decomposition $L = K_L A_L^+ K_L$ or $L = K_L Z_L^+ K_L$ with $K_L \subset K$, with $\mathbf{A}_L \subset \mathbf{A}$, and with $A_L^+ \cap A^+$ non-compact. Let $\mu_L : L \rightarrow E_L^+$ be the corresponding Cartan projection. We naturally see E_L as a line in E . If $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 1, then E_L^+ is a half-line in E^+ ; if $\mathbf{L}$ has semisimple $\mathbf{k}$ -rank 0, then $E_L^+ = E_L$ is a line in E , intersecting E^+ in a half-line, and

$$\mu(g) = E^+ \cap W \cdot \mu_L(g)$$

for all $g \in L$. Since all norms on E are equivalent, we may assume that $\|\cdot\|$ is the W -invariant Euclidean norm introduced in Section 2. By composing μ_L with some isomorphism from E_L to $\mathbb{R}$, we get a Cartan projection $\mu_L^{\mathbb{R}} : L \rightarrow \mathbb{R}$ with

$$|\mu_L^{\mathbb{R}}(g)| = \|\mu(g)\|$$

for all $g \in L$. For every $\alpha \in \Delta$ there are constants $t_\alpha^+, t_\alpha^- \geq 0$ such that

$$(5.1) \quad \langle \alpha, \mu(g) \rangle = \begin{cases} t_\alpha^+ |\mu_L^{\mathbb{R}}(g)| & \text{if } \mu_L^{\mathbb{R}}(g) \geq 0, \\ t_\alpha^- |\mu_L^{\mathbb{R}}(g)| & \text{if } \mu_L^{\mathbb{R}}(g) \leq 0. \end{cases}$$

Let $\Delta_L = \{\alpha \in \Delta, t_\alpha^\pm > 0\}$ denote the set of simple roots of $\mathbf{A}$ in $\mathbf{G}$ whose restriction to $\mathbf{A}_L$ is nontrivial. Let E_{Δ_L} denote the subspace of E spanned by the coroots $\check{\alpha}$ for $\alpha \in \Delta_L$, and let $\text{pr}_{E_{\Delta_L}} : E \rightarrow E_{\Delta_L}$ denote the orthogonal projection on E_{Δ_L} . Then

$$|v|_{E_{\Delta_L}} = \|\text{pr}_{E_{\Delta_L}}(v)\|$$

defines a seminorm $|\cdot|_{E_{\Delta_L}}$ on E . For $\alpha \in \Delta$, let χ_α denote the highest weight of the representation (ρ_α, V_α) of $\mathbf{G}$ introduced in Subsection 2.5. Recall that $\langle \chi_\alpha, \check{\alpha} \rangle \neq 0$ and $\langle \chi_\alpha, \check{\beta} \rangle = 0$ for all $\beta \in \Delta \setminus \{\alpha\}$, hence $\{\langle \chi_\alpha, \cdot \rangle, \alpha \in \Delta_L\}$ is a basis of the dual of E_{Δ_L} . Thus the function

$$v \mapsto \sum_{\alpha \in \Delta_L} |\langle \chi_\alpha, v \rangle|$$

is a norm on E_{Δ_L} . Since all norms on E_{Δ_L} are equivalent, there exists $c \geq 1$ such that

$$(5.2) \quad c^{-1} \cdot \sum_{\alpha \in \Delta_L} |\langle \chi_\alpha, v \rangle| \leq |v|_{E_{\Delta_L}} \leq c \cdot \sum_{\alpha \in \Delta_L} |\langle \chi_\alpha, v \rangle|$$

for all $v \in E$.

5.2. Norm of the projection on E_{Δ_L} . The main step in the proof of Proposition 5.1 consists of the following proposition, which gives an upper bound for the seminorm $|\cdot|_{E_{\Delta_L}}$.

Proposition 5.2. *Under the assumptions of Proposition 5.1, for any $\delta > 0$ there exist a finite subset F'_δ of Γ , a neighborhood $\mathcal{U}'_\delta \subset \text{Hom}(\Gamma, G)$ of the natural inclusion, and a constant $C'_\delta \geq 0$ such that any $\gamma \in \Gamma$ may be written as $\gamma = \gamma_0 \dots \gamma_n$ for some $\gamma_0, \dots, \gamma_n \in F'_\delta$ with*

- (1) $n \leq \delta \sum_{i=0}^n \|\mu(\gamma_i)\|$,
- (2) $\sum_{i=1}^n \|\mu(\gamma_i)\| = \|\sum_{i=1}^n \mu(\gamma_i)\|$,
- (3) $\|\mu(\varphi(\gamma_i)) - \mu(\gamma_i)\| \leq 1$ for all $\varphi \in \mathcal{U}'_\delta$ and $0 \leq i \leq n$,
- (4) for all $\varphi \in \mathcal{U}'_\delta$,

$$\left| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right|_{E_{\Delta_L}} \leq \delta \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + C'_\delta.$$

To prove Proposition 5.2, we use Propositions 3.2 and 4.1, together with Lemma 4.4.

Proof. Let $D > 0$ be the constant and $\mathcal{C}_L$ the compact subset of N_L^- given by Proposition 4.1. By Lemma 4.4, for any $\alpha \in \Delta_L$, the set $\mathcal{C}_L$ is contained in $\text{Int}(\mathcal{C}_\alpha)P_\alpha$ for some compact subset $\mathcal{C}_\alpha$ of N_α^- , where $\text{Int}(\mathcal{C}_\alpha)$ denotes the interior of $\mathcal{C}_\alpha$. Let $r_\alpha, R_\alpha > 0$ be the corresponding constants given by Proposition 3.2. Fix $\delta > 0$ and choose $R > D$ large enough so that $\frac{1}{R-D} \leq \delta$ and $\min(t_\alpha^+, t_\alpha^-)(R-D)-1 \geq R_\alpha$ for all $\alpha \in \Delta_L$, where t_α^+ and t_α^- are defined by (5.1). Let F'_δ be the set of elements $\gamma \in \Gamma$ such that $|\mu_L^\mathbb{R}(\gamma)| \leq R+D$, and F''_δ the subset of elements $\gamma \in F'_\delta$ such that $|\mu_L^\mathbb{R}(\gamma)| \geq R-D$. Note that F'_δ et F''_δ are finite since $\mu_L^\mathbb{R}$ is a proper map and Γ is discrete in L . Let $\mathcal{U}'_\delta \subset \text{Hom}(\Gamma, G)$ be the neighborhood of the natural inclusion whose elements φ satisfy the following two conditions:

- $\|\mu(\varphi(\gamma)) - \mu(\gamma)\| \leq 1$ and $|\langle \alpha, \mu(\varphi(\gamma)) - \mu(\gamma) \rangle| \leq 1$ for all $\gamma \in F'_\delta$ and all $\alpha \in \Delta_L$,
- $\ell_{\varphi(\gamma)} k_{\varphi(\gamma')} \in \mathcal{C}_\alpha P_\alpha$ for all $\gamma, \gamma' \in F''_\delta$ with $\ell_\gamma k_{\gamma'} \in \mathcal{C}_L P_L$ and all $\alpha \in \Delta_L$,

where for $g \in G$ we write $g = k_g z_g \ell_g$ with $k_g, \ell_g \in K$ and $z_g \in Z^+$. We claim that F'_δ and $\mathcal{U}'_\delta$ satisfy the conclusions of Proposition 5.2 for some constant C'_δ . Indeed, let $\gamma \in \Gamma$. By Proposition 4.1, we may write $\gamma = \gamma_0 \dots \gamma_n$ for some elements $\gamma_0 \in F'_\delta$ and $\gamma_1, \dots, \gamma_n \in F''_\delta$ such that

- $\ell_{\gamma_i} k_{\gamma_{i+1}} \in \mathcal{C}_L P_L$ for all $1 \leq i \leq n-1$,
- $\mu_L^\mathbb{R}(\gamma_1), \dots, \mu_L^\mathbb{R}(\gamma_n)$ are all ≥ 0 or all ≤ 0 .

This last condition implies that $\mu_L^\mathbb{R}(\gamma_1), \dots, \mu_L^\mathbb{R}(\gamma_n)$ all belong to the same half-line in E^+ , hence

$$\sum_{i=1}^n \|\mu(\gamma_i)\| = \left\| \sum_{i=1}^n \mu(\gamma_i) \right\|.$$

Moreover, since $\gamma_1, \dots, \gamma_n \in F''_\delta$, we have

$$n \leq \frac{1}{R-D} \cdot \sum_{i=0}^n |\mu_L^\mathbb{R}(\gamma_i)| \leq \delta \sum_{i=0}^n \|\mu(\gamma_i)\|.$$

Let $\varphi \in \mathcal{U}'_\delta$. According to (5.2), in order to prove Condition (4) it is sufficient to bound

$$\left| \left\langle \chi_\alpha, \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right\rangle \right|$$

for all $\alpha \in \Delta_L$. For $\alpha \in \Delta_L$ and $1 \leq i \leq n$ we have

$$\begin{aligned} \langle \alpha, \mu(\varphi(\gamma_i)) \rangle &\geq \langle \alpha, \mu(\gamma_i) \rangle - 1 \\ &\geq \min(t_\alpha^+, t_\alpha^-) |\mu_L^{\mathbb{R}}(\gamma_i)| - 1 \\ &\geq \min(t_\alpha^+, t_\alpha^-)(R - D) - 1 \geq R_\alpha \end{aligned}$$

and $\ell_{\varphi(\gamma_i)} k_{\varphi(\gamma_{i+1})} \in \mathcal{C}_\alpha P_\alpha$. Proposition 3.2 thus implies that

$$\begin{aligned} \left| \left\langle \chi_\alpha, \mu(\varphi(\gamma_1 \dots \gamma_n)) - \sum_{i=1}^n \mu(\varphi(\gamma_i)) \right\rangle \right| &\leq n r_\alpha \\ &\leq \frac{r_\alpha}{R - D} \sum_{i=0}^n \|\mu(\gamma_i)\|. \end{aligned}$$

On the other hand, by (5.2) and (2.4),

$$\begin{aligned} \left| \left\langle \chi_\alpha, \mu(\varphi(\gamma)) - \mu(\varphi(\gamma_1 \dots \gamma_n)) \right\rangle \right| &\leq c \|\mu(\varphi(\gamma_0))\| \\ &\leq c \cdot \max_{f \in F'_\delta} (\|\mu(f)\| + 1). \end{aligned}$$

By the triangular inequality, we finally get

$$\left| \left\langle \chi_\alpha, \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right\rangle \right| \leq \frac{r_\alpha}{R - D} \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + C'_\delta$$

with

$$C'_\delta = c \cdot \max_{f \in F'_\delta} (\|\mu(f)\| + |\langle \chi_\alpha, \mu(f) \rangle| + 2).$$

By (5.2), this implies Condition (4) whenever

$$\sum_{\alpha \in \Delta_L} \frac{c r_\alpha}{R - D} \leq \delta,$$

which holds for R large enough. $\square$

5.3. Proof of Proposition 5.1. Proposition 5.1 follows from Proposition 5.2 and from the following general observation.

Lemma 5.3. *Let $(E, \|\cdot\|)$ be a Euclidean space, with corresponding scalar product $\langle \cdot, \cdot \rangle$, and E_1 a subspace of E . For any $x \in E$, let $|x|_1 = \|\text{pr}_1(x)\|$, where $\text{pr}_1 : E \rightarrow E_1$ denotes the orthogonal projection on E_1 . For any $\delta > 0$, fix $C''_\delta \geq 0$ and let $\mathcal{I}_\delta$ be the set of pairs $(x, x') \in E^2$ such that*

- (1) $x \in E_1$,
- (2) $|x' - x|_1 \leq 2\delta \|x\| + C''_\delta$,
- (3) $\|x'\| \leq (1 + \delta) \|x\| + C''_\delta$.

Then

$$\sup_{(x, x') \in \mathcal{I}_\delta, x \neq 0} \frac{\|x' - x\| - 4C''_\delta/\delta}{\|x\|} \xrightarrow{\delta \rightarrow 0} 0.$$

Proof. Let $\delta \in]0, 1]$ and $(x, x') \in \mathcal{I}_\delta$. If $\|x\| \leq C''_\delta/\delta$, then

$$\|x' - x\| \leq \|x'\| + \|x\| \leq (2 + \delta) \|x\| + C''_\delta \leq \frac{4C''_\delta}{\delta}.$$

If $\|x\| \geq C''_\delta/\delta$, then

$$\|x'\| \leq (1 + 2\delta) \|x\|$$

and

$$\langle x, x' \rangle = \langle x, \text{pr}_1(x') \rangle \geq \|x\|^2 - \|x\| \|x - x'\|_1 \geq (1 - 3\delta) \|x\|^2.$$

To conclude, note that for $y \in E$ with $\|y\| = 1$, the diameter of the set

$$\{y' \in E, \quad 1 - 3\delta \leq \langle y, y' \rangle \leq \|y'\| \leq 1 + 2\delta\}$$

uniformly tends to 0 with δ . $\square$

Proof of Proposition 5.1. Fix $\varepsilon \in]0, 1]$. For $\delta \in]0, 1]$, let F'_δ , $\mathcal{U}'_\delta$, and C'_δ be given by Proposition 5.2. Let $\gamma \in \Gamma$. By Proposition 5.2, we may write $\gamma = \gamma_0 \dots \gamma_n$ for some $\gamma_0, \dots, \gamma_n \in F'_\delta$ such that

- (1) $n \leq \delta \sum_{i=0}^n \|\mu(\gamma_i)\|$,
- (2) $\sum_{i=1}^n \|\mu(\gamma_i)\| = \|\sum_{i=1}^n \mu(\gamma_i)\|$,
- (3) $\|\mu(\varphi(\gamma_i)) - \mu(\gamma_i)\| \leq 1$ for all $\varphi \in \mathcal{U}'_\delta$ and $0 \leq i \leq n$,
- (4) for all $\varphi \in \mathcal{U}'_\delta$,

$$\left| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right|_{E_{\Delta_L}} \leq \delta \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + C'_\delta.$$

Conditions (1), (3), and (4), together with (2.3) and the triangular inequality, imply that for all $\varphi \in \mathcal{U}'_\delta$,

$$\begin{aligned} \left| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\gamma_i) \right|_{E_{\Delta_L}} &\leq \left| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right|_{E_{\Delta_L}} + \sum_{i=0}^n \|\mu(\varphi(\gamma_i)) - \mu(\gamma_i)\| \\ &\leq 2\delta \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + C'_\delta + 1 \end{aligned}$$

and

$$\|\mu(\varphi(\gamma))\| \leq \sum_{i=0}^n \|\mu(\varphi(\gamma_i))\| \leq (1 + \delta) \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + 1.$$

Moreover, Condition (2) implies that

$$(5.3) \quad \sum_{i=0}^n \|\mu(\gamma_i)\| \leq \left\| \sum_{i=0}^n \mu(\gamma_i) \right\| + 2 \max_{g \in F'_\delta} \|\mu(g)\|.$$

Therefore, for $\varphi \in \mathcal{U}'_\delta$, Lemma 5.3 applies to

$$\begin{aligned} E_1 &= E_{\Delta_L}, \\ C''_\delta &= C'_\delta + 1 + 4 \max_{g \in F'_\delta} \|\mu(g)\|, \end{aligned}$$

$$\begin{aligned} x &= \sum_{i=0}^n \mu(\gamma_i), \\ x' &= \mu(\varphi(\gamma)) : \end{aligned}$$

we obtain that if δ is small enough, then

$$(5.4) \quad \left\| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\gamma_i) \right\| \leq \frac{\varepsilon}{4} \left\| \sum_{i=0}^n \mu(\gamma_i) \right\| + \frac{4C''_\delta}{\delta}$$

for all $\varphi \in \mathcal{U}'_\delta$. Now Conditions (1) and (3), together with the triangular inequality and (5.3), imply that

$$\begin{aligned} \left\| \left(\sum_{i=0}^n \mu(\varphi(\gamma_i)) \right) - \left(\sum_{i=0}^n \mu(\gamma_i) \right) \right\| &\leq \delta \left(\sum_{i=0}^n \|\mu(\gamma_i)\| \right) + 1, \\ &\leq \delta \left\| \sum_{i=0}^n \mu(\gamma_i) \right\| + 2\delta \max_{g \in F'_\delta} \|\mu(g)\| + 1. \end{aligned}$$

Therefore, if δ is small enough, then

$$\left\| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right\| \leq \frac{\varepsilon}{2} \left\| \sum_{i=0}^n \mu(\gamma_i) \right\| + C''_\delta$$

for all $\varphi \in \mathcal{U}'_\delta$, where

$$C''_\delta = \frac{4C''_\delta}{\delta} + 2\delta \max_{g \in F'_\delta} \|\mu(g)\| + 1.$$

In particular, taking φ to be the natural inclusion of Γ in G , we get

$$(5.5) \quad \left\| \sum_{i=0}^n \mu(\gamma_i) \right\| \leq \frac{1}{1 - \frac{\varepsilon}{2}} (\|\mu(\gamma)\| + C''_\delta) \leq 2\|\mu(\gamma)\| + 2C''_\delta,$$

hence

$$\left\| \mu(\varphi(\gamma)) - \sum_{i=0}^n \mu(\varphi(\gamma_i)) \right\| \leq \varepsilon \|\mu(\gamma)\| + 3C''_\delta$$

for all $\varphi \in \mathcal{U}'_\delta$ whenever δ is small enough. Finally, Condition (1), together with (5.3) and (5.5), implies that

$$n \leq 2\delta \|\mu(\gamma)\| + 3C''_\delta.$$

Thus the triple $(F_\varepsilon, \mathcal{U}_\varepsilon, C_\varepsilon) = (F'_\delta, \mathcal{U}'_\delta, 3C''_\delta)$ satisfies the conclusions of Proposition 5.1 for δ small enough. $\square$

5.4. Properness and deformation. Let us now briefly explain how to deduce Theorems 1.1 and 1.3 from Theorem 1.4.

Theorem 1.3 follows from Theorem 1.4 and from the *properness criterion* of Benoist ([Ben], Cor. 5.2) and Kobayashi ([Ko2], Th. 1.1). Under the assumptions of Theorem 1.3, this criterion states that a subgroup Γ of G acts properly on G/H if and only if the set $\mu(\Gamma) \cap (\mu(H) + \mathcal{C})$ is bounded for any compact subset $\mathcal{C}$ of E . This condition means that the set $\mu(\Gamma)$ “gets away from $\mu(H)$ at infinity”.

Proof of Theorem 1.3. We may assume that $\mathbf{G}$, $\mathbf{H}$, and $\mathbf{L}$ are all connected. Let $G = KA^+K$ or $G = KZ^+K$ be a Cartan decomposition of G , and let $\mu : G \rightarrow E^+$ be the corresponding Cartan projection. Endow E with a W -invariant norm $\|\cdot\|$ as in Section 2. By Remark 4.3, we may assume that L admits a Cartan decomposition $L = K_L A_L^+ K_L$ or $L = K_L Z_L^+ K_L$ with $K_L \subset K$ and $\mathbf{A}_L \subset \mathbf{A}$. If $\mu_L : L \rightarrow E_L^+$ denotes the corresponding Cartan projection, then E_L is naturally seen as a line in E and

$$\mu(\ell) = E^+ \cap (W \cdot \mu_L(\ell))$$

for all $\ell \in L$. Thus $\mu(L)$ is contained in the union U_L of two half-lines of E^+ . Using Remark 4.3 again, there is an element $g \in G$ such that gHg^{-1} admits a Cartan decomposition $gHg^{-1} = K_H A_H^+ K_H$ or $gHg^{-1} = K_H Z_H^+ K_H$ with $K_H \subset K$ and $\mathbf{A}_H \subset \mathbf{A}$. The set $\mu(gHg^{-1})$ is contained in a finite union U_H of subspaces of E intersected with E^+ , parametrized by the Weyl group W . By (2.4), the Hausdorff distance between $\mu(gHg^{-1})$ and $\mu(H)$ is $\leq 2\|\mu(g)\|$. Therefore $U_L \cap U_H = \{0\}$ by the properness criterion, and there is a constant $\varepsilon > 0$ such that

$$d(\mu(\ell), \mu(H)) \geq 2\varepsilon \|\mu(\ell)\| - 2\|\mu(g)\|$$

for all $\ell \in L$. By Theorem 1.4, there is a neighborhood $\mathcal{U}_\varepsilon \subset \text{Hom}(\Gamma, G)$ of the natural inclusion and a constant $C_\varepsilon \geq 0$ such that

$$\|\mu(\varphi(\gamma)) - \mu(\gamma)\| \leq \varepsilon \|\mu(\gamma)\| + C_\varepsilon$$

for all $\varphi \in \mathcal{U}_\varepsilon$ and $\gamma \in \Gamma$. Fix $\varphi \in \mathcal{U}_\varepsilon$. For all $\gamma \in \Gamma$,

$$\begin{aligned} d(\mu(\varphi(\gamma)), \mu(H)) &\geq d(\mu(\gamma), \mu(H)) - \|\mu(\varphi(\gamma)) - \mu(\gamma)\| \\ &\geq \varepsilon \|\mu(\gamma)\| - C_\varepsilon - 2\|\mu(g)\|. \end{aligned}$$

Therefore, using the fact that Γ is discrete in G and μ is a proper map, we get that $\mu(\varphi(\Gamma)) \cap (\mu(H) + \mathcal{C})$ is finite for any compact subset $\mathcal{C}$ of E . By the properness criterion, this implies that $\varphi(\Gamma)$ acts properly on G/H . It also implies that $\varphi(\Gamma)$ is discrete in G and that the kernel of φ is finite. Since Γ is torsion-free, φ is injective. $\square$

Proof of Theorem 1.1. By Theorem 1.3, there is a neighborhood $\mathcal{U} \subset \text{Hom}(\Gamma, G)$ of the natural inclusion such that any $\varphi \in \mathcal{U}$ is injective and $\varphi(\Gamma)$ is discrete in G , acting properly discontinuously on G/H . Since $\varphi \in \mathcal{U}$ is injective, $\varphi(\Gamma)$ has the same cohomological dimension as Γ . We conclude using the fact, due to Kobayashi ([Ko1], Cor. 5.5), that a torsion-free discrete subgroup of G acts cocompactly sur G/H if and only if its cohomological dimension is $d(G) - d(H)$, where $d(G)$ (resp. $d(H)$) denotes the dimension of the symmetric space of G (resp. of H). $\square$

6. APPLICATION TO THE COMPACT QUOTIENTS OF $\text{SO}(2n, 2)/\text{U}(n, 1)$

Fix an integer $n \geq 1$. Note that $\text{U}(n, 1)$ naturally embeds in $\text{SO}(2n, 2)$ by identifying the Hermitian form $|z_1|^2 + \dots + |z_n|^2 - |z_{n+1}|^2$ on $\mathbb{C}^{n+1}$ with the quadratic form $x_1^2 + \dots + x_{2n}^2 - x_{2n+1}^2 - x_{2n+2}^2$ on $\mathbb{R}^{2n+2}$. As Kulkarni [Kul] pointed out, $\text{U}(n, 1)$, seen as a subgroup of $\text{SO}(2n, 2)$, acts transitively on the anti-de Sitter space

$$\begin{aligned} \text{AdS}^{2n+1} &= \{(x_1, \dots, x_{2n+2}) \in \mathbb{R}^{2n+2}, x_1^2 + \dots + x_{2n}^2 - x_{2n+1}^2 - x_{2n+2}^2 = -1\} \\ &\simeq \text{SO}(2n, 2)/\text{SO}(2n, 1). \end{aligned}$$

The stabilizer of $(0, \dots, 0, 1)$ is the compact subgroup $\text{U}(n)$, hence AdS^{2n+1} identifies with $\text{U}(n, 1)/\text{U}(n)$ and the action of $\text{U}(n, 1)$ on $\text{SO}(2n, 2)/\text{SO}(2n, 1)$ is proper. By duality, the action of $\text{SO}(2n, 1)$ on $\text{SO}(2n, 2)/\text{U}(n, 1)$ is proper and transitive. In particular, any uniform lattice Γ of $\text{SO}(2n, 1)$ provides a standard compact quotient $\Gamma \backslash \text{SO}(2n, 2)/\text{U}(n, 1)$ of $\text{SO}(2n, 2)/\text{U}(n, 1)$.

Corollary 1.2 follows from Theorem 1.1 and from the existence of Zariski-dense deformations in $\text{SO}(m, 2)$ of certain uniform lattices of $\text{SO}(m, 1)$. Such

deformations can be obtained by a bending construction due to Johnson and Millson. This construction is presented in [JM] for deformations in $\mathrm{SO}(m+1, 1)$ or in $\mathrm{PGL}_{m+1}(\mathbb{R})$, but not in $\mathrm{SO}(m, 2)$. For the reader's convenience, we shall describe Johnson and Millson's construction in the latter case, and check that the deformations obtained in this way are indeed Zariski-dense in $\mathrm{SO}(m, 2)$.

From now on we use Gothic letters to denote the Lie algebras of real Lie groups (*e.g.* $\mathfrak{g}$ for G).

6.1. Uniform arithmetic lattices of $\mathrm{SO}(m, 1)$. Fix $m \geq 2$. The uniform lattices of $\mathrm{SO}(m, 1)$ considered by Johnson and Millson are obtained in the following classical way. Fix a square-free integer $r \geq 2$ and identify $\mathrm{SO}(m, 1)$ with the special orthogonal group of the quadratic form

$$x_1^2 + \dots + x_m^2 - \sqrt{r}x_{m+1}^2$$

on $\mathbb{R}^{m+1}$. Let $\mathcal{O}_r$ denote the ring of integers of the quadratic field $\mathbb{Q}(\sqrt{r})$. The group $\Gamma = \mathrm{SO}(m, 1) \cap M_{m+1}(\mathcal{O}_r)$ is a uniform lattice in $\mathrm{SO}(m, 1)$ (see [Bo1] for instance). For any ideal I of $\mathcal{O}_r$, the congruence subgroup $\Gamma \cap (1 + M_{m+1}(I))$ has finite index in Γ , hence is a uniform lattice in $\mathrm{SO}(m, 1)$. By [MR], after replacing Γ by such a congruence subgroup, we may assume that it is torsion-free. Then $M = \Gamma \backslash \mathbb{H}^m$ is a m -dimensional compact hyperbolic manifold whose fundamental group identifies with Γ . By [JM], Lem. 7.1 & Th. 7.2, after possibly replacing Γ again by some congruence subgroup, we may assume that $N = \Gamma_0 \backslash \mathbb{H}^{m-1}$ is a connected, orientable, totally geodesic hypersurface of M , where

$$\Gamma_0 = \Gamma \cap \mathrm{SO}(m-1, 1)$$

and where

$$\mathbb{H}^{m-1} \simeq \{(x_2, \dots, x_{m+1}) \in \mathbb{R}^m, x_2^2 + \dots + x_m^2 - \sqrt{r}x_{m+1}^2 = -1 \text{ and } x_{m+1} > 0\}$$

is embedded in

$$\mathbb{H}^m \simeq \{(x_1, \dots, x_{m+1}) \in \mathbb{R}^{m+1}, x_1^2 + \dots + x_m^2 - \sqrt{r}x_{m+1}^2 = -1 \text{ and } x_{m+1} > 0\}$$

in the natural way. Since the centralizer of Γ_0 in $\mathrm{SO}(m, 2)$ contains a subgroup isomorphic to $\mathrm{SO}(1, 1) \simeq \mathbb{R}^*$, the idea of the bending construction is to deform Γ “along this centralizer”, as we shall now explain.

6.2. Deformations in the separating case. Assume that N separates M into two components M_1 and M_2 , and let Γ_1 (resp. Γ_2) denote the fundamental group of M_1 (resp. of M_2). By van Kampen's theorem, Γ is the amalgamated product $\Gamma_1 *_{\Gamma_0} \Gamma_2$. Fix an element $Y \in \mathfrak{so}(m, 2) \setminus \mathfrak{so}(m, 1)$ that belongs to the Lie algebra of the centralizer of Γ_0 in $\mathrm{SO}(m, 2)$. Following Johnson and Millson, we consider the deformations of Γ in $\mathrm{SO}(m, 2)$ that are given, for $t \in \mathbb{R}$, by

$$\varphi_t(\gamma) = \begin{cases} \gamma & \text{for } \gamma \in \Gamma_1, \\ e^{tY} \gamma e^{-tY} & \text{for } \gamma \in \Gamma_2. \end{cases}$$

Note that $\varphi_t : \Gamma \rightarrow \mathrm{SO}(m, 2)$ is well defined since e^{tY} centralizes Γ_0 . Moreover, it is injective and $\varphi_t(\Gamma)$ is discrete in $\mathrm{SO}(m, 2)$. We now check Zariski-density.

Lemma 6.1. *For $t \neq 0$ small enough, $\varphi_t(\Gamma)$ is Zariski-dense in $\mathrm{SO}(m, 2)$.*

We need the following remark.

Remark 6.2. For $m \geq 2$, the only Lie subalgebra of $\mathfrak{so}(m, 2)$ that strictly contains $\mathfrak{so}(m, 1)$ is $\mathfrak{so}(m, 2)$.

Indeed, $\mathfrak{so}(m, 2)$ decomposes uniquely into a direct sum $\mathfrak{so}(m, 1) \oplus \mathbb{R}^{m+1}$ of irreducible $\mathrm{SO}(m, 1)$ -modules, where $\mathrm{SO}(m, 1)$ acts on $\mathfrak{so}(m, 1)$ (resp. on $\mathbb{R}^{m+1}$) by the adjoint (resp. natural) action.

Proof of Lemma 6.1. Recall that $\mathrm{SO}(m, 2)$ is Zariski-connected. Therefore, in order to prove that $\varphi_t(\Gamma)$ is Zariski-dense in $\mathrm{SO}(m, 2)$, it is sufficient to prove that the Lie algebra of $\overline{\varphi_t(\Gamma)}$ is $\mathfrak{so}(m, 2)$, where $\overline{\varphi_t(\Gamma)}$ denotes the Zariski closure of $\varphi_t(\Gamma)$ in $\mathrm{SO}(m, 2)$.

By [JM], Lem. 5.9, the groups Γ_1 and Γ_2 are Zariski-dense in $\mathrm{SO}(m, 1)$. By [JM], Cor. 5.3, and [Ser], § I.5.2, Cor. 1, they naturally embed in Γ . Therefore $\overline{\varphi_t(\Gamma)}$ contains both $\mathrm{SO}(m, 1)$ and $e^{tY}\mathrm{SO}(m, 1)e^{-tY}$, and the Lie algebra of $\overline{\varphi_t(\Gamma)}$ contains both $\mathfrak{so}(m, 1)$ and the Lie algebra of $e^{tY}\mathrm{SO}(m, 1)e^{-tY}$. By Remark 6.2, in order to prove that $\varphi_t(\Gamma)$ is Zariski-dense in $\mathrm{SO}(m, 2)$, it is sufficient to prove that the Lie algebra of $e^{tY}\mathrm{SO}(m, 1)e^{-tY}$ is not $\mathfrak{so}(m, 1)$.

But if the Lie algebra of $e^{tY}\mathrm{SO}(m, 1)e^{-tY}$ were $\mathfrak{so}(m, 1)$, then we would have $e^{tY}\mathrm{SO}(m, 1)^\circ e^{-tY} = \mathrm{SO}(m, 1)^\circ$, i.e., e^{tY} would belong to the normalizer $N_{\mathrm{SO}(m, 2)}(\mathrm{SO}(m, 1)^\circ)$ of the identity component $\mathrm{SO}(m, 1)^\circ$ of $\mathrm{SO}(m, 1)$. Recall that the exponential map induces a diffeomorphism between a neighborhood $\mathcal{U}$ of 0 in $\mathfrak{so}(m, 2)$ and a neighborhood $\mathcal{V}$ of 1 in $\mathrm{SO}(m, 2)$, which itself induces a one-to-one correspondence between $\mathcal{U} \cap \mathfrak{n}_{\mathrm{so}(m, 2)}(\mathfrak{so}(m, 1))$ and $\mathcal{V} \cap N_{\mathrm{SO}(m, 2)}(\mathrm{SO}(m, 1)^\circ)$. Therefore, if we had $e^{tY} \in N_{\mathrm{SO}(m, 2)}(\mathrm{SO}(m, 1)^\circ)$ for some $t \neq 0$ small enough, then we would have

$$Y \in \mathfrak{n}_{\mathrm{so}(m, 2)}(\mathfrak{so}(m, 1)) = \{X \in \mathfrak{so}(m, 2), \mathrm{ad}(X)(\mathfrak{so}(m, 1)) = \mathfrak{so}(m, 1)\}.$$

But Remark 6.2 implies that $\mathfrak{n}_{\mathrm{so}(m, 2)}(\mathfrak{so}(m, 1))$ is equal to $\mathfrak{so}(m, 1)$, since it contains $\mathfrak{so}(m, 1)$ and is different from $\mathfrak{so}(m, 2)$. Thus we would have $Y \in \mathfrak{so}(m, 1)$, which would contradict our choice of Y . $\square$

6.3. Deformations in the nonseparating case. We now assume that $S = M \setminus N$ is connected. Let $j_1 : \Gamma_0 \rightarrow \pi_1(S)$ and $j_2 : \Gamma_0 \rightarrow \pi_1(S)$ denote the inclusions in $\pi_1(S)$ of the fundamental groups of the two sides of N . The group Γ is a HNN extension of $\pi_1(S)$, i.e., it is generated by $\pi_1(S)$ and by some element $\nu \in \Gamma$ such that

$$\nu j_1(\gamma) \nu^{-1} = j_2(\gamma)$$

for all $\gamma \in \Gamma_0$. Fix an element $Y \in \mathfrak{so}(m, 2) \setminus \mathfrak{so}(m, 1)$ that belongs to the Lie algebra of the centralizer of $j_1(\Gamma_0)$ in $\mathrm{SO}(m, 2)$. Following Johnson and Millson, we consider the deformations of Γ in $\mathrm{SO}(m, 2)$ that are given, for $t \in \mathbb{R}$, by

$$\begin{cases} \varphi_t(\gamma) = \gamma & \text{for } \gamma \in \pi_1(S), \\ \varphi_t(\nu) = \nu e^{tY}. \end{cases}$$

Note that $\varphi_t : \Gamma \rightarrow \mathrm{SO}(m, 2)$ is well defined since e^{tY} centralizes $j_1(\Gamma_0)$. Moreover, it is injective and $\varphi_t(\Gamma)$ is discrete in $\mathrm{SO}(m, 2)$.

Lemma 6.3. *For $t \neq 0$ small enough, $\varphi_t(\Gamma)$ is Zariski-dense in $\mathrm{SO}(m, 2)$.*

Proof. Let $\overline{\varphi_t(\Gamma)}$ denote the Zariski closure of $\varphi_t(\Gamma)$ in $\mathrm{SO}(m, 2)$. By [JM], Lem. 5.9, the group $\pi_1(S)$ is Zariski-dense in $\mathrm{SO}(m, 1)$, hence $\overline{\varphi_t(\Gamma)}$ contains both $\mathrm{SO}(m, 1)$ and νe^{tY} . But $\nu \in \mathrm{SO}(m, 1)$, hence $e^{tY} \in \overline{\varphi_t(\Gamma)}$. Therefore $\overline{\varphi_t(\Gamma)}$ contains both $\mathrm{SO}(m, 1)$ and $e^{tY}\mathrm{SO}(m, 1)e^{-tY}$, and we may conclude as in the proof of Lemma 6.1. $\square$

REFERENCES

- [Bas] H. BASS, *Covering theory for graphs of groups*, J. Pure Appl. Algebra 89 (1993), p. 3–47.
- [Ben] Y. BENOIST, *Actions propres sur les espaces homogènes réductifs*, Ann. Math. 144 (1996), p. 315–347, announced in C. R. Acad. Sci. Paris, Ser. I, Math. 319 (1994), p. 937–940.
- [Bo1] A. BOREL, *Compact Clifford-Klein forms of symmetric spaces*, Topology 2 (1963), p. 111–122.
- [Bo2] A. BOREL, *Linear algebraic groups*, Second edition, Graduate Texts in Mathematics 126, Springer-Verlag, New York, 1991.
- [BoT] A. BOREL, J. TITS, *Groupes réductifs*, Publ. Math. Inst. Hautes Études Sci. 27 (1965), p. 55–150.
- [Bou] M. BOURDON, *Structure conforme au bord et flot géodésique d'un $CAT(-1)$ -espace*, Enseign. Math. 41 (1995), p. 63–102.
- [BH] M. R. BRIDSON, A. HAEFLIGER, *Metric spaces of non-positive curvature*, Grundlehren der mathematischen Wissenschaften 319, Springer-Verlag, Berlin, 1999.
- [BT1] F. BRUHAT, J. TITS, *Groupes réductifs sur un corps local : I. Données radicielles valuées*, Publ. Math. Inst. Hautes Études Sci. 41 (1972), p. 5–251.
- [BT2] F. BRUHAT, J. TITS, *Groupes réductifs sur un corps local : II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Publ. Math. Inst. Hautes Études Sci. 60 (1984), p. 5–184.
- [Che] C. CHEVALLEY, *Théorie des groupes de Lie : II. Groupes algébriques*, Actualités Sci. Ind. 1152, Hermann & Cie, Paris, 1951.
- [Ghy] É. GHYS, *Déformations des structures complexes sur les espaces homogènes de $\mathrm{SL}_2(\mathbb{C})$* , J. Reine Angew. Math. 468 (1995), p. 113–138.
- [Gol] W. M. GOLDMAN, *Nonstandard Lorentz space forms*, J. Differ. Geom. 21 (1985), p. 301–308.
- [Gui] O. GUICHARD, *Groupes plongés quasi-isométriquement dans un groupe de Lie*, Math. Ann. 330 (2004), p. 331–351.
- [Hel] S. HELGASON, *Differential geometry, Lie groups, and symmetric spaces*, Corrected reprint of the 1978 original, Graduate Studies in Mathematics 34, American Mathematical Society, Providence, RI, 2001.
- [JM] D. JOHNSON, J. J. MILLSON, *Deformation spaces associated to compact hyperbolic manifolds*, in *Discrete groups in geometry and analysis*, p. 48–106, Prog. Math. 67, Birkhäuser, Boston, MA, 1987.
- [Kar] F. I. KARPELEVICH, *Surfaces of transitivity of semisimple groups of motions of a symmetric space*, Soviet Math. Dokl. 93 (1953), p. 401–404.
- [Kas] F. KASSEL, *Proper actions on corank-one reductive homogeneous spaces*, J. Lie Theory 18 (2008), p. 961–978.
- [Kli] B. KLINGLER, *Complétude des variétés lorentziennes à courbure constante*, Math. Ann. 306 (1996), p. 353–370.

- [Ko1] T. KOBAYASHI, *Proper action on a homogeneous space of reductive type*, Math. Ann. 285 (1989), p. 249–263.
- [Ko2] T. KOBAYASHI, *Criterion for proper actions on homogeneous spaces of reductive groups*, J. Lie Theory 6 (1996), p. 147–163.
- [Ko3] T. KOBAYASHI, *Deformation of compact Clifford-Klein forms of indefinite-Riemannian homogeneous manifolds*, Math. Ann. 310 (1998), p. 394–408.
- [KY] T. KOBAYASHI, T. YOSHINO, *Compact Clifford-Klein forms of symmetric spaces — revisited*, Pure and Applied Mathematics Quaterly 1 (2005), p. 591–653.
- [Kul] R. S. KULKARNI, *Proper actions and pseudo-Riemannian space forms*, Adv. Math. 40 (1981), p. 10–51.
- [Lan] E. LANDVOGT, *Some functorial properties of the Bruhat-Tits building*, J. Reine Angew. Math. 518 (2000), p. 213–241.
- [Mar] G. A. MARGULIS, *Discrete subgroups of semisimple Lie groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) 17, Springer-Verlag, Berlin, 1991.
- [MR] J. J. MILLSON, M. S. RAGHUNATHAN, *Geometric construction of cohomology for arithmetic groups I*, in *Geometry and analysis*, p. 103–123, Indian Acad. Sci., Bangalore, 1980.
- [Mos] G. D. MOSTOW, *Some new decomposition theorems for semi-simple groups*, Mem. Amer. Math. Soc. 14 (1955), p. 31–54.
- [Qui] J.-F. QUINT, *Cônes limites des sous-groupes discrets des groupes réductifs sur un corps local*, Transform. Groups 7 (2002), p. 247–266.
- [Rag] M. S. RAGHUNATHAN, *On the first cohomology of discrete subgroups of semisimple Lie groups*, Amer. J. Math. 87 (1965), p. 103–139.
- [Rou] G. ROUSSEAU, *Euclidean buildings*, in “Nonpositive curvature geometries, discrete groups and rigidity”, Proceedings of the 2004 Grenoble summer school, Séminaires et Congrès 18, Société Mathématique de France, Paris, 2008.
- [Sal] F. SALEIN, *Variétés anti-de Sitter de dimension 3 exotiques*, Ann. Inst. Fourier 50 (2000), p. 257–284.
- [Sel] A. SELBERG, *On discontinuous groups in higher-dimensional symmetric spaces*, in “Collected papers”, vol. 1, p. 475–492, Springer-Verlag, Berlin, 1989.
- [Ser] J.-P. SERRE, *Arbres, amalgames, $SL(2)$* , Astérisque 46, Société Mathématique de France, Paris, 1977.
- [Tit] J. TITS, *Représentations linéaires irréductibles d’un groupe réductif sur un corps quelconque*, J. Reine Angew. Math. 247 (1971), p. 196–220.
- [Wei] A. WEIL, *Remarks on the cohomology of groups*, Ann. Math. 80 (1964), p. 149–157.
- [Zeg] A. ZEGHIB, *On closed anti-de Sitter spacetimes*, Math. Ann. 310 (1998), p. 695–716.

DÉPARTEMENT DE MATHÉMATIQUES, BÂTIMENT 425, FACULTÉ DES SCIENCES D’ORSAY,
 UNIVERSITÉ PARIS-SUD 11, 91405 ORSAY CEDEX, FRANCE
E-mail address: fanny.kassel@math.u-psud.fr